



Kingdom of Cambodia

Trends in Maternal and Child Health in Cambodia, 2000-2014

Further Analysis of the 2000, 2005, 2010, and 2014
Cambodia Demographic and Health Surveys



DHS Further Analysis Reports No. 106

Kingdom of Cambodia



Trends in Maternal and Child Health in Cambodia, 2000-2014

Further Analysis of the 2000, 2005, 2010, and 2014 Cambodia Demographic and Health Surveys

DHS Further Analysis Reports No. 106

Lindsay Mallick
Courtney Allen
Rathavuth Hong

ICF
Rockville, Maryland, USA

January 2018



Acknowledgments:

The authors would like to thank Shireen Assaf and Tom Pullum for their Stata coding support.

Editor: Bryant Robey

Document Production: Joan Wardell

This report presents findings from a further analysis study undertaken as part of the follow-up to the 2014 Cambodia Demographic and Health Survey (CDHS). ICF provided technical assistance for the project. This report is a publication of the DHS program, which is designed to collect, analyze, and disseminate data on fertility, family planning, maternal and child health, nutrition, and HIV/AIDS. Funding was provided by the U.S. Agency for International Development (USAID) through the DHS Program (#AID-OAA-C-13-00095). The opinions expressed herein are those of the authors and do not necessarily reflect the views of USAID and other cooperating agencies.

The 2014 Cambodia Demographic and Health Survey (2014 CDHS) is part of The DHS Program, a worldwide project that assists countries in the collection of data to monitor and evaluate population, health, and nutrition programs. Funding was provided by the Royal Government of Cambodia (RGC), the United States Agency for International Development (USAID), the Australian Department of Foreign Affairs and Trade (Australia-DFAT), the United Nations Population Fund (UNFPA), the United Nations Children's Fund (UNICEF), the Japan International Cooperation Agency (JICA), the Korean International Cooperation Agency (KOICA), and the Health Sector Support Program—Second Phase (HSSP-2).

Additional information about the survey can be obtained from the National Institute of Statistics; 386 Monivong Boulevard, Sangkat Boeng Keng Kang 1, Chamkar Mon, Phnom Penh, Cambodia; Telephone: (855) 23-213650; E-mail: ssythan@hotmail.com; Internet: www.nis.gov.kh; and the Directorate General for Health, Ministry of Health, 80 Samdech Penn North Boulevard (289), Sangkat Boeungkak 2, Tuol Kork, Phnom Penh, Cambodia; Telephone: (855) 23-885970/23-884909; E-mail: webmaster@moh.gov.kh; Internet: www.moh.gov.kh.

Additional information about The DHS Program can be obtained from ICF, 530 Gaither Road, Suite 500, Rockville, MD 20850 USA; Telephone: 301-572-0200, Fax: 301-572-0999, E-mail: info@DHSprogram.com, Internet: www.DHSprogram.com.

Suggested citation:

Mallick, Lindsay, Courtney Allen, and Rathavuth Hong. 2016. *Trends in Maternal and Child Health in Cambodia, 2000-2014. Further Analysis of the 2000, 2005, 2010, and 2014 Cambodia Demographic and Health Surveys*. DHS Further Analysis Reports No. 106. Rockville, Maryland, USA: ICF

CONTENTS

TABLES	v
FIGURES	vii
ABSTRACT	ix
1 INTRODUCTION	1
2 DATA AND METHODS	3
2.1 Data	3
2.2 Measures.....	3
2.3 Analysis	5
3 RESULTS	7
3.1 Maternal Health.....	7
3.1.1 Antenatal Care	7
3.1.2 Components of Antenatal Care.....	10
3.1.3 Delivery Care Indicators	12
3.1.4 Postnatal Care	14
3.1.5 The Continuum of Maternal Health Care.....	16
3.2 Child Health.....	17
3.2.1 Size at Birth	18
3.2.2 Early Initiation of Breastfeeding.....	21
3.2.3 Full Immunization.....	23
3.2.4 Care Seeking for Diarrhea	25
3.2.5 Treatment for Diarrhea	27
3.2.6 Nutritional Indices	29
4 DISCUSSION	33
4.1 Maternal Health.....	33
4.2 Child Health.....	34
4.3 Conclusion.....	35
APPENDIX	37
REFERENCES	55

TABLES

Table 1.	Description of the Cambodia Demographic and Health Surveys (CDHS) included in the analysis	3
Table 2.	Maternal health indicators included in the analysis	4
Table 3.	Child health indicators included in the analysis.....	5
Table 4.	Distribution of women age 15-49 with a live birth in the 5 years preceding the survey, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS.....	7
Table 5.	Distribution of children under age 5, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS.....	18
Appendix Table 1.	National-level trends for maternal health indicators, Cambodia 2000, 2005, 2010, and 2014 DHS	39
Appendix Table 2.	Percentage of women with four or more antenatal care visits for their most recent pregnancy among women age 15-49 with a live birth in the 5 years preceding the survey, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	40
Appendix Table 3.	Among women age 15-49 with a live birth in the 5 years preceding the survey who received antenatal care for their most recent birth, percentage who were informed of signs of complications, Cambodia 2000, 2005, 2010, and 2014 DHS	41
Appendix Table 4.	Percentage of births that were assisted by a skilled birth attendant, among all births in the 5 years preceding the survey to women age 15-49, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	42
Appendix Table 5.	Percentage of women who received a postnatal check-up within 2 days of delivering their most recent child among women age 15-49 with a live birth in the 2 years preceding the survey, according to background characteristics, Cambodia 2005, 2010, and 2014 DHS.....	43
Appendix Table 6.	Distribution of women with antenatal care (four or more visits), a skilled birth attendant at delivery, or postnatal check-up within 2 days of delivering their most recent child among women age 15-49 with a live birth in the 2 years preceding the survey, according to background characteristics, Cambodia 2005, 2010, and 2014 DHS	44
Appendix Table 7.	National-level trends for child health indicators, Cambodia 2000, 2005, 2010, and 2014 DHS.....	45
Appendix Table 8.	Percentage of children born in the last 5 years who were smaller than average by mother's estimation, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	46
Appendix Table 9.	Percentage of last-born children in the 2 years preceding the survey who started breastfeeding within 1 hour of birth, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	48
Appendix Table 10.	Percentage of children age 12-23 months who are fully immunized, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	49

Appendix Table 11.	Among children under age 5 who experienced diarrhea in the 2 weeks before the survey, percentage for whom treatment was sought from a health facility or provider, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	50
Appendix Table 12.	Among children under age 5 who experienced diarrhea in the 2 weeks before the survey, percentage given ORT or increased fluids, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	51
Appendix Table 13.	Among children under age 5, percentage who are stunted, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	52

FIGURES

Figure 1.	Infant and under-5 mortality rates Cambodia 2000, 2005, 2010, and 2014 DHS.....	1
Figure 2.	Maternal mortality ratio, Cambodia 2000, 2005, 2010, and 2014 DHS	2
Figure 3.	Indicators of antenatal care for the most recent birth of women with a live birth in the 5 years preceding the survey, Cambodia 2000, 2005, 2010, and 2014 DHS.....	8
Figure 4.	Percentage of women with four or more antenatal care visits for their most recent pregnancy among women age 15-49 with a live birth in the 5 years preceding the survey, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	9
Figure 5.	Indicators of components of antenatal care for the most recent birth of women with a live birth in the 5 years preceding the survey among women who received antenatal care, Cambodia 2000, 2005, 2010, and 2014 DHS	10
Figure 6.	Among women age 15-49 with a live birth in the 5 years preceding the survey who received antenatal care for their most recent birth, the percentage who were informed of signs of complications, Cambodia 2000, 2005, 2010, and 2014 DHS.....	11
Figure 7.	Indicators of delivery care for all births in the 5 years preceding the survey, Cambodia 2000, 2005, 2010, and 2014 DHS.....	12
Figure 8.	Percentage of births that were assisted by a skilled birth attendant, among all births in the 5 years preceding the survey to women age 15-49, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	13
Figure 9.	Percentage of women who received a postnatal check-up within 2 days of delivering their most recent child among women age 15-49 with a live birth in the 2 years preceding the survey, according to background characteristics, Cambodia 2005, 2010, and 2014 DHS.....	15
Figure 10.	Among women with a live birth in the 2 years before the survey, the percent distribution of the maternal health services received during the most recent birth and pregnancy, Cambodia 2005, 2010, and 2014 DHS	17
Figure 11.	Percentage of children born in the last 5 years who were not weighed, Cambodia 2000, 2005, 2010, and 2014 DHS	19
Figure 12.	Percentage of live births in the last 5 years who were estimated to be smaller than average by mothers, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	20
Figure 13.	Percentage of last-born children in the 2 years preceding the survey who started breastfeeding within 1 hour of birth, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS.....	22
Figure 14.	Percentage of children age 12-23 months who received each vaccine, Cambodia 2000, 2005, 2010, and 2014 DHS.....	23
Figure 15.	Percentage of children age 12-23 months who are fully immunized, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS	24

Figure 16.	Among children under age 5 who experienced diarrhea in the 2 weeks before the survey, percentage for whom treatment was sought from a health facility or provider, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS.....	26
Figure 17.	Among children under age 5 who experienced diarrhea in the 2 weeks before the survey, percentage given ORT or increased fluids, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS.....	28
Figure 18.	Among children under age 5, percentage who are stunted, wasted, or underweight, Cambodia 2000, 2005, 2010, and 2014 DHS.....	30
Figure 19.	Among children under age 5, percentage who are stunted, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS.....	30


ABSTRACT

This report examines the trends in maternal and child health over a 14-year period (2000-2014) in Cambodia. The analysis draws from four nationally representative household surveys conducted by The Demographic and Health Surveys Program in 2000, 2005, 2010, and 2014. We examined estimates of indicators of antenatal care, delivery, and postnatal care for women as well as size at birth, breastfeeding, immunization coverage, care-seeking, and nutritional status of children. We tested for the significance of the change of these estimates at the national level as well as by sociodemographic characteristics. The results revealed significant improvement in a number of indicators, although the greatest changes appeared between 2000 and 2010, with fewer gains seen between 2010 and 2014. For many indicators examined in this report, significant disparities continued over time, most notably among regions, wealth quintiles, and levels of mothers' education. Overall, the trends in maternal and child health care in Cambodia are positive, though continued efforts are needed to ensure that there is equitable access to health services for all women and children.

1 INTRODUCTION

Over the past 2 decades, The Demographic and Health Surveys (DHS) Program has conducted four nationally representative surveys in Cambodia to assess the health of the population, including maternal and child health. These surveys indicate that reproductive health outcomes in Cambodia have improved substantially since 2000. By the end of 2014, Cambodia had achieved several of the health-related Millennium Development Goals (MDGs). For example, as Figure 1 shows, from 2000 to 2014 the decrease in both the infant mortality rate and under-5 mortality rate far exceeded the MDG target set for 2015, to reduce under-5 mortality by two-thirds from 1990 to 2015 (National Institute of Public Health/Cambodia, National Institute of Statistics/Cambodia, and ORC Macro 2006; National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ICF International 2015; National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ICF Macro 2011; National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ORC Macro 2001).

Figure 1. Infant and under-5 mortality rates Cambodia 2000, 2005, 2010, and 2014 DHS




Note: Mortality ratios obtained from The DHS Program STATcompiler: <https://www.statcompiler.com>.

Similarly, the maternal mortality ratio (MMR) declined significantly between 2000 and 2014. Figure 2 presents MMR for the 7 years preceding each survey, expressed per 100,000 live births, calculated as the age-adjusted maternal mortality rate times 100 divided by the age-adjusted general fertility rate. Between the first two surveys (2000 and 2005) in which the DHS measured MMR, the trend was discouraging. Subsequently, efforts were made to reduce pregnancy-related deaths and to halt and reverse the trend. After 2005, MMR began to improve, eventually reaching the MDG target by 2014, to reduce maternal mortality by three-quarters from 1990 to 2015.

Cambodia has experienced rapid economic growth in the last 2 decades. Through dedicated government resources, as well as international donor funding, access to good-quality health care has improved with the implementation of health equity funds, vouchers, and community-based health insurance schemes (Dingle, Goodman, and Powell-Jackson 2013; Van de Poel et al. 2014; WHO n.d.). Much of the improvement in childhood and maternal morbidity and mortality may be attributed to the general increase in provision and use of maternal and child health services, while other socioeconomic indicators such as

universal primary education and eradication of extreme poverty have also shown substantial improvement. Nonetheless, Cambodia’s health sector still faces many challenges, such as disparities and inequities within socioeconomic sub-groups (Dingle, Goodman, and Powell-Jackson 2013) and lack of improvement in some health indicators. For example, while many efforts have been made to reduce the proportion of children who are undernourished, the prevalence of stunting and underweight among children under age 5 has remained high, below the MDG targets set for 2015 (Rikimaru 2015).

Figure 2. Maternal mortality ratio, Cambodia 2000, 2005, 2010, and 2014 DHS



Note: Maternal mortality ratios obtained from the The DHS Program STATcompiler: <https://www.statcompiler.com>. Point estimate includes the lower and upper bounds of the 95% confidence intervals.

The objective of the present analysis is to describe trends in several key maternal and child health indicators from 2000 to 2014, using data from the Cambodia Demographic and Health Surveys (CDHS) Surveys. This report addresses the prevailing maternal and child health situation in Cambodia and the trends in antenatal care (ANC), delivery, postnatal care (PNC), birth size, and breastfeeding, as well as immunization, care and treatment of illness, and nutritional status among children. The report also examines the interventions and practices that may contribute to reducing maternal and child health mortality. The primary objective is to provide information for policymakers and program administrators to help assess the current situation and to design future maternal and child health programs.

2 DATA AND METHODS

2.1 Data

This analysis used data from the four CDHS surveys in 2000, 2005, 2010, and 2014. Data collected by the CDHS surveys are comparable over time because the surveys use the same sampling design, model questionnaires, data collection techniques, measures, and methods of analysis. The 2000, 2010, and 2014 surveys were implemented by the Cambodian National Institute of Statistics (NIS) of the Ministry of Planning and the Directorate General for Health (DGH) of the Ministry of Health. The 2005 survey was carried out by the NIS and the National Institute of Public Health and Research (NIPH) of the Ministry of Health. Table 1 presents information on dates of fieldwork and sample sizes for the households and women interviewed in the CDHS.

Table 1. Description of the Cambodia Demographic and Health Surveys (CDHS) included in the analysis

Year	Date of fieldwork	Implementing organization	Number of households interviewed	Number of women aged 15-49 interviewed
2000	February-July 2000	NIS ¹ & DGH ²	12,236	15,351
2005	September 2005-March 2006	NIS & NIPH ³	14,243	16,823
2010	July 2010-January 2011	NIS & DGH	15,667	18,754
2014	June-December 2014	NIS & DGH	15,825	17,578

¹ NIS: National Institute of Statistics (Ministry of Planning); ² DGH: Directorate General for Health (Ministry of Health); ³ NPH: National Institute of Public Health and Research (Ministry of Health)

For each survey in Cambodia, the DHS used a complex, two-stage sample design first to select clusters and then to select households resulting in a nationally representative sample. The interviewers collected data from members of the selected households, including women age 15-49, about their health attitudes, behavior, and outcomes. If a woman had a live birth in the 5 years preceding the survey, her survey included questions about the care she received during pregnancy, birth, and in the postnatal period. Additional questions also asked about the health of and care for her children. This study examines women with a live birth in the 5 years preceding each survey and the children born during those 5 years. This includes 5,714 women and 8,715 children in the 2000 CDHS, 5,865 women and 7,789 children in 2005, 6,472 women and 8,200 children in 2010, and 5,973 women and 7,253 children in 2014.

2.2 Measures

The indicators assessed in this report provide a snapshot of the health and the care received among mothers during and after pregnancy and birth, and for their children up to age 5. However, the definitions of some of these key indicators changed over the survey period. In order to ensure comparability across surveys, the definitions of some variables are standardized. For example, earlier surveys only assessed PNC among women who delivered at home, versus all women in more recent surveys. Additionally, in more recent surveys this indicator is only presented for women with a birth in the 2 years preceding the survey. Therefore, we do not analyze PNC for women interviewed in the 2000 CDHS. We present results only among women with a birth in the most recent 2 years preceding the survey (versus 5 years). Among the child health indicators, we examine trends in oral rehydration therapy (ORT) given to children with diarrhea symptoms in the 2 weeks preceding the survey. In the earliest survey, this definition included oral rehydration solution (ORS), recommended home fluids (RHF), rice water, or increased fluids. For all four

surveys, in order to compare them over time, we defined ORT as receiving ORS or increased fluids. Additionally, where initial breastfeeding is calculated, the indicator is for the last-born child in the 2 years preceding the survey. This is different from the denominator in earlier surveys, which included all children born in the 5 years preceding the survey in 2000 and only last-born children in the 5 years preceding the survey in 2005. Table 2 presents a list of the maternal health indicators, the standard definition used to calculate the indicators across each survey, and the sample size for each. Table 3 shows the respective information for the child health indicators.

Table 2. Maternal health indicators included in the analysis

Indicator	Definition	Population Base	Sample Size			
			2000	2005	2010	2014
Four or more antenatal care visits (ANC)	Percentage of women with four or more antenatal care visits for their most recent pregnancy	Women age 15-49 with a live birth in the 5 years preceding the survey	5,714	5,865	6,472	5,973
Timing of first ANC	Percentage of women who received ANC in the first 4 months of pregnancy	Women age 15-49 with a live birth in the 5 years preceding the survey	5,714	5,865	6,472	5,973
Mother took iron syrup/tablets during pregnancy	Percentage of women who were given iron syrup/tablets during their most recent pregnancy	Women age 15-49 with a live birth in the 5 years preceding the survey	5,714	5,865	6,472	5,973
Mother took deworming medication during pregnancy	Percentage of women who consumed deworming medication during their most recent pregnancy	Women age 15-49 with a live birth in the 5 years preceding the survey	5,714	5,865	6,472	5,973
Blood pressure checked during ANC	Percentage of women who had their blood pressure checked during an ANC visit during their most recent pregnancy	Women age 15-49 with a live birth in the 5 years preceding the survey who had at least one ANC visit	2,543	4,213	5,804	5,704
Informed of pregnancy complications during ANC	Percentage of women who were informed of pregnancy complications during an ANC visit during their most recent pregnancy	Women age 15-49 with a live birth in the 5 years preceding the survey who had at least one ANC visit	2,543	4,213	5,804	5,704
Birth delivered in a facility	Percentage of births that were delivered in a facility	Children born in the 5 years preceding the survey	8,715	7,789	8,200	7,253
Births assisted by a skilled birth attendant (SBA)	Percentage of births that were assisted by an SBA	Children born in the 5 years preceding the survey	8,715	7,789	8,200	7,253
Births delivered by Caesarean section	Percentage of births that were delivered by caesarean section	Children born in the 5 years preceding the survey	8,715	7,789	8,200	7,253
Postnatal care for the mother	Percentage of women who received a postnatal check-up within 2 days of delivering their most recent birth	Women age 15-49 with a live birth in the 2 years preceding the survey	n/a	3,083	3,187	2,944
Continuum of care	Percentage of women who received any combination of ANC (4 or more visits), skilled birth attendant, and postnatal care	Women age 15-49 with a live birth in the 2 years preceding the survey	n/a	3,083	3,187	2,944

Table 3. Child health indicators included in the analysis

Indicator	Definition	Population Base	Sample Size			
			2000	2005	2010	2014
Mother's estimate of baby's size at birth	Percent distribution of live births in the 5 years preceding the survey by mother's estimate of baby's size at birth	Children under age 5	8,715	7,789	8,200	7,253
Early initiation of breastfeeding	Among last-born children who were born in the 2 years preceding the survey, the percentage who started breastfeeding within 1 hour	Most recent birth in the 2 years preceding the survey	7,825	5,711	3,187	2,944
Full immunization	Percentage of children age 12-23 months who received specific vaccines at any time before the survey (according to a vaccination card or the mother's report)	Children age 12-23 months	1,253	1,517	1,614	1,460
Care seeking for diarrhea from a facility or health provider	Among children under age 5 who had diarrhea in the 2 weeks preceding the survey, the percentage for whom advice or treatment was sought from a health facility or provider	Children under age 5 with diarrhea in the 2 weeks preceding the survey	1,385	1,420	1,161	902
ORT or increased fluids as part of treatment in children who had diarrhea	Among children under age 5 who had diarrhea in the 2 weeks preceding the survey, the percentage given ORT or increased fluids	Children under age 5 with diarrhea in the 2 weeks preceding the survey	1,385	1,420	1,161	902
Nutritional status: stunting	Percentage of children under age 5 who are at least 2 standard deviations below the median for the reference population for height-for-age	Children under age 5 with anthropometric assessments	3,372	3,587	3,975	4,893

2.3 Analysis

We used data from the four CDHS surveys to investigate the changes in indicators of maternal and child health over time. We conducted tests of association to identify significant changes between each survey, as well as between the first survey (2000 CDHS) and the most recent survey (2014 CDHS). For selected indicators, we performed additional tests of associations to assess the significance of relationships between these indicators and sociodemographic and health covariates within each survey and across surveys. These variables included place of residence (urban or rural), region, wealth quintile, and mother's education (none, primary, or secondary and higher). We grouped the 19 provinces of Cambodia into five regions: Phnom Penh, Plain, Great Lake, Coastal, and Plateau¹. Depending on the indicator, we also included maternal age at birth, parity or birth order, birth interval, ANC, facility delivery, assistance at birth, whether the mother smokes cigarettes, age of the child, and sex of the child. We used Stata 15 for all calculations, weighting our estimates and adjusting our analysis for the complex sample design using the *svy* command.

This report presents the results of the trends in indicators of maternal and child health in line graphs, which demonstrate the percentage at each time point, the change between successive surveys and between the first and the last survey, and the significance of the change between surveys. In each figure, a solid line between 2 time points represents a significant change from one survey to the next, and a dotted line indicates no significant change. Significant differences between the first survey (2000 CDHS) and the last survey

¹ The Phnom Penh region includes only the capital city, Phnom Penh. The Plain region includes the provinces of Kampong Cham, Kandal, Prey Veng, Svay Rieng, and Takeo. The Great Lake region includes the provinces of Banteay Meanchey, Battambang, Pailin, Kampong Chhnang, Kampong Thom, Poursat, and Siem Reap. The Coastal region includes the provinces of Kampot, Koh Kong, Kep, and Preah Sihanouk. The Plateau region includes the provinces of Kampong Speu, Kratie, Mondol Kiri, Preah Vihear, Rattanak Kiri, Stung Treng, and Otdor Mean Chey.

(2014 CDHS) are marked with asterisks in the legend for each indicator or subgroup. The number of asterisks seen denotes the p-value: * is a p-value <0.05, ** is <0.01, and *** is <0.001. No asterisks indicate no significant changes.

3 RESULTS

3.1 Maternal Health

Table 4 presents the background characteristics of women with a live birth in the 5 years preceding each survey. In all four surveys the majority of these women were rural residents and about two in every five lived in the Plain region. The majority in each survey had a primary education, though the percentage of women with secondary or higher education increased over the survey period. Among these women, most were age 20-34 and had three children or fewer. The proportion of women with four, five, or six or more children decreased over the survey period. Appendix Table 1 contains the national-level trends and the results of the significance tests of changes over time.

Table 4. Distribution of women age 15-49 with a live birth in the 5 years preceding the survey, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000		2005		2010		2014	
	%	N	%	N	%	N	%	N
Place of residence								
Urban	13.6	779	14.1	827	16.2	1,050	14.7	876
Rural	86.4	4,935	85.9	5,039	83.8	5,421	85.3	5,096
Region								
Phnom Penh	5.9	336	8.1	476	8.3	538	9.0	535
Plain	41.9	2,393	39.7	2,329	40.0	2,587	36.7	2,193
Great Lake	31.8	1,819	31.0	1,816	30.3	1,958	29.9	1,785
Coastal	7.9	451	7.4	437	6.8	440	6.3	378
Plateau	12.5	715	13.8	807	14.7	948	18.1	1,081
Wealth quintile								
Lowest	25.1	1,436	25.2	1,477	24.5	1,585	22.8	1,359
Second	22.2	1,269	22.5	1,320	21.3	1,380	20.3	1,215
Middle	20.2	1,152	18.4	1,077	19.0	1,229	19.0	1,133
Fourth	18.3	1,043	17.1	1,003	17.9	1,155	17.9	1,069
Highest	14.3	814	16.9	988	17.4	1,123	20.0	1,196
Education								
None	32.0	1,827	23.1	1,356	17.5	1,133	13.5	805
Primary	53.7	3,069	59.4	3,482	56.2	3,635	51.9	3,100
Secondary+	14.3	818	17.5	1,028	26.3	1,703	34.6	2,068
Mother's age at birth								
<20	8.1	465	9.2	540	8.6	555	10.4	620
20-34	68.6	3,921	70.2	4,118	76.0	4,917	79.5	4,749
35-49	23.3	1,329	20.6	1,206	15.4	999	10.1	603
Parity								
1	17.1	975	24.4	1,430	30.6	1,980	35.3	2,109
2-3	35.9	2,053	40.5	2,378	45.3	2,931	47.4	2,828
4-5	22.7	1,296	20.5	1,200	15.5	1,001	12.7	760
6+	24.3	1,391	14.6	857	8.7	560	4.6	276
Total	100.0	5,714	100.0	5,865	100.0	6,472	100.0	5,973


3.1.1 Antenatal Care

Figure 3 shows the national-level percentage of women in each survey with a live birth who obtained key ANC services during their most recent pregnancy. Specifically, this figure shows the trends in receiving four or more ANC visits, receiving ANC in the first three months of pregnancy, consumption of iron supplementation, and consumption of intestinal parasite medication. The 2000 survey did not collect data on consumption of parasite medication.

Overall, the trend is positive for each indicator. There were significant improvements for each indicator at the national level between successive rounds of surveys and also between the first and last

surveys. The levels and trends are similar for receipt of four or more ANC visits and for early initiation of ANC. Results on both indicators show that about 10% of women received coverage in 2000, rising to almost 80% in 2014. Consumption of intestinal parasite medication also rose about 70 percentage points between 2005 and 2014. By 2014, nearly 100% of women with a pregnancy in the 5 years preceding the survey consumed iron supplements, compared with 20% in 2000.


Figure 3. Indicators of antenatal care for the most recent birth of women with a live birth in the 5 years preceding the survey, Cambodia 2000, 2005, 2010, and 2014 DHS



Note: A solid line indicates a significant change between two surveys. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: *<0.05, **<0.01, ***<0.001.

Figure 4 shows the percentage of women making four or more ANC visits in their last pregnancy, by background characteristics, including place of residence, region, wealth, education, age at birth, and parity. Appendix Table 2 includes the corresponding percentages, confidence intervals, differences, and significance levels, as well as the significance of the difference for each background characteristic. Between the first and the last surveys, each demographic group showed significant improvement. Between each of the surveys, each category of women also significantly improved, with the exception of women in Phnom Penh and women with six or more births between 2010 and 2014. Each demographic group demonstrated significant differences by receipt of four or more ANC visits for each survey. The disparities were smallest between urban and rural residence and by age at birth. For region, wealth, and education, the disparities between the lowest and highest categories were approximately 30 percentage points. By region, women in Phnom Penh were the most likely to make four more ANC visits while the lowest coverage appeared to be women in the Plateau region. The gap by parity widened over time. Women with any children similarly obtained the recommended four or more visits in 2000 (between 5% and 15%) regardless of their number of births. The proportion of women who made the recommended number of ANC visits increased more over the survey period among lower parity women compared with higher parity women. By 2014, less than 40% of women with six or more births made four or more ANC visits, compared with over 80% of primiparous women.

Figure 4. Percentage of women with four or more antenatal care visits for their most recent pregnancy among women age 15-49 with a live birth in the 5 years preceding the survey, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS




Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * < 0.05, ** < 0.01, *** < 0.001.

3.1.2 Components of Antenatal Care

Figure 5 shows the trend in selected components of ANC for women who had any ANC visits during their most recent pregnancy resulting in a live birth in the 5 years preceding each survey. Coverage of both indicators—blood pressure measurement and being informed of signs of complications—significantly increased from 2000 to 2014, by 35 percentage points and 42 percentage points respectively. The greatest change for both indicators was between 2000 and 2014, with only a small change between 2010 and 2014; the increase in the percentage of women informed of complications is not significant between 2010 and 2014. Figure 6 disaggregates this indicator by women’s background characteristics.

Figure 5. Indicators of components of antenatal care for the most recent birth of women with a live birth in the 5 years preceding the survey among women who received antenatal care, Cambodia 2000, 2005, 2010, and 2014 DHS




Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: *<0.05, **<0.01, ***<0.001.

Figure 6 shows that for almost all categories, the change between 2010 and 2014 is also not significant, while the overall trend between the first and the last surveys is highly significant. The trend over time is unique by region. In Phnom Penh there was a sharp increase (23 percentage points) from 2005 to 2010, but a significant decrease of 7 percentage points from 2010 to 2014, with a net increase of 24 percentage points from 2000 to 2014. In the Great Lake region, the percentage of women who were informed of signs of pregnancy complications during ANC increased by 50 percentage points from 2000 to 2014.

The disparities within each background characteristic have also narrowed over time. In 2000, significant differences existed between demographic categories. In contrast, in 2014, significant differences appear by category only for education, age at child’s birth, and parity. This suggests that, once women obtain care, providers do not give unequal treatment by place of residence, region, and wealth. Appendix Table 3 contains supplementary information.

Figure 6. Among women age 15-49 with a live birth in the 5 years preceding the survey who received antenatal care for their most recent birth, the percentage who were informed of signs of complications, Cambodia 2000, 2005, 2010, and 2014 DHS




Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * < 0.05, ** < 0.01, *** < 0.001.

3.1.3 Delivery Care Indicators

Figure 7 demonstrates the trend in delivery care received for all births in the 5 years preceding each survey, from 2000 to 2014. All of the indicators of care at birth, including delivery in a health facility, skilled birth attendance (SBA), and caesarian section (C-section), demonstrate significant increases between each survey, and between the first and the last surveys. Facility delivery increased by 73 percentage points, to 83% in 2014, and SBA increased by 57 percentage points, to almost 90% in 2014. The biggest change in these two indicators occurred between 2005 and 2010. Few births were delivered by C-section; the proportion increased from 1% in 2000 to 3% in 2014.


Figure 7. Indicators of delivery care for all births in the 5 years preceding the survey, Cambodia 2000, 2005, 2010, and 2014 DHS




Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: *<0.05, **<0.01, ***<0.001.

Figure 8 presents, for each survey, the proportion of births in the preceding 5 years delivered by a skilled birth attendant, for each background characteristic of women. In addition to the demographic characteristics for the birth, including place of residence, region, wealth, education, maternal age at birth, and birth order, we present the proportions of births by number of ANC visits (for the most recent birth only) and whether or not the birth occurred in a health facility. (See Appendix Table 4 for additional information.)

Figure 8. Percentage of births that were assisted by a skilled birth attendant, among all births in the 5 years preceding the survey to women age 15-49, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS





Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * <0.05 , ** <0.01 , *** <0.001 . Births with don't know or missing responses are included with the category "0-3 visits" for ANC and the "home, other" category for place of delivery.

For most characteristics, the largest changes occurred between 2005 and 2010. Among demographic characteristics, including place of residence, region, wealth, and education, the disparities diminished over time. For example, in 2000 the difference in having a skilled attendant at birth ranged from 15% in the lowest wealth quintile to 81% in the highest quintile—over 60 percentage points. With substantial improvements among women in the lower wealth quintiles, the gap between lowest and highest declined to just over 20 percentage points by 2014. However, the differences within groups remain significant, with a p-value <0.001 for each demographic characteristic.


This analysis also presents the proportion of births delivered by a skilled birth attendant (SBA) by the number of ANC visits and whether the delivery was in a health facility or not. The disaggregation by ANC includes only the most recent birth to the interviewed mother. Births to women who made four or more ANC visits during their pregnancy were significantly more likely to be delivered by an SBA, though by 2014 having an SBA was common for both groups. In all four surveys, births were almost universally delivered by an SBA if they were delivered in a health facility, while less than 40% of home births had an SBA present. Among home births, however, the proportion assisted by an SBA increased significantly over the survey period.


3.1.4 Postnatal Care

Figure 9, complemented by Appendix Table 5, displays the proportion of women who received a postnatal check-up (PNC) within 2 days of delivery among women with a birth in the 2 years preceding each survey. Data from the 2000 CDHS are not comparable and thus are not included. The proportion of women who received PNC increased significantly, from 65% in 2005 to 90% in 2014; however, the rate of change was not consistent across sociodemographic indicators. For example, the percentage of women with PNC in the Plateau region increased significantly, by 27 percentage points from 2005 to 2014, though almost none of this change occurred between 2005 and 2010. There was no significant change among women in the highest wealth quintile between 2010 and 2014. By region, wealth, and education, the disparities diminished over time. By 2014 there were no significant differences by maternal age at birth and

only marginally significant differences by parity. The most notable disparity was by skilled attendance at birth: 94% of women with skilled attendance at birth had PNC, whereas only 51% of women without a birth attendant received PNC. Further, the proportion of women receiving PNC among women who did not have an SBA did not significantly increase from 2005 to 2014.

Figure 9. Percentage of women who received a postnatal check-up within 2 days of delivering their most recent child among women age 15-49 with a live birth in the 2 years preceding the survey, according to background characteristics, Cambodia 2005, 2010, and 2014 DHS






Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2005 and 2014 is indicated in the legend with asterisks to represent the p-value: * <0.05 , ** <0.01 , *** <0.001 . Women with don't know or missing information are included with the category "0-3 visits" for ANC and the "home, other" category for attendance at birth.

3.1.5 The Continuum of Maternal Health Care

The continuum of care refers to three key services a woman can receive from pregnancy to birth and in the postpartum period. Ensuring that women receive maternal health interventions throughout this cycle is a critical strategy for reducing maternal and child mortality and morbidity (Kerber et al. 2007). The first service, ANC, may be the time a woman begins using health care regularly; ANC visits can help women learn about the importance of receiving delivery and postnatal services. Use of ANC has a positive effect on continued use of maternal services (Adjiwanou and LeGrand 2013; Wang and Hong 2015). Figure 10 shows the proportion of women with a birth in the 2 years preceding each of three Cambodia surveys (2005, 2010, and 2014) who made four or more ANC visits at last pregnancy, had an SBA present at delivery, and received PNC within 2 days after delivery. This figure clearly shows not only that the proportion of women receiving each indicator of maternal health service has increased but also that the proportion of women receiving the continuum of all three services has increased. In 2014, 69% of women received all three services compared with just 18% in 2005—a three-fold increase in nine years. Moreover, the proportion of women receiving none of the three maternal services decreased, from 23% in 2005 to 2% in 2014. The proportion of women who did not receive the recommended four or more ANC visits but who had an SBA at delivery and received PNC remained near 20% over the survey period. Appendix Table 6 includes the percentages for each category and the significance of change for each category over time.

Figure 10. Among women with a live birth in the 2 years before the survey, the percent distribution of the maternal health services received during the most recent birth and pregnancy, Cambodia 2005, 2010, and 2014 DHS



3.2 Child Health

Table 5 presents the background characteristics of children under age 5 who were born to women interviewed in the Women’s Questionnaire. In all four surveys, most children lived in rural areas and the most common region of residence was the Plain region. The majority of children were born to mothers with primary education, women age 20-34, and women with three children or fewer. Appendix Table 7 shows the national-level trends and the results of the significance tests of changes over time. Appendix Tables 8-13 contain the indicator percentages and confidence intervals over the four surveys, as well as level of significance for each sociodemographic variable and each measured difference between two consecutive surveys and the first and last surveys that correspond to each indicator, disaggregated by women’s background characteristics.

Table 5. Distribution of children under age 5, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS


	2000		2005		2010		2014	
	%	N	%	N	%	N	%	N
Age of child¹								
<6 mo	11.9	882	10.7	782	9.3	729	10.6	750
6-11 mo	10.7	788	10.6	773	10.6	826	10.8	761
1 years	16.9	1,253	20.8	1,517	20.6	1,614	20.7	1,460
2 years	18.7	1,379	19.4	1,418	20.6	1,610	19.4	1,368
3 years	20.8	1,541	19.6	1,430	19.6	1,537	19.0	1,343
4 years	21.0	1,553	19.0	1,389	19.3	1,514	19.5	1,376
Sex of child								
Male	50.9	4,161	50.9	3,901	50.1	4,245	51.8	3,636
Female	49.1	4,014	49.1	3,887	49.9	3,955	48.2	3,617
Place of residence								
Urban	13.2	1,076	14.0	1,093	15.6	1,281	14.4	1,041
Rural	86.8	7,098	86.0	6,696	84.4	6,919	85.7	6,212
Region								
Pnom Penh	5.3	433	7.9	614	7.9	647	8.6	626
Plain	41.2	3,368	38.2	2,974	39.4	3,232	36.9	2,677
Great Lakes	33.3	2,719	31.7	2,467	31.0	2,542	30.0	2,177
Coastal	7.5	614	7.6	593	6.7	546	6.1	444
Plateau	12.7	1,041	14.7	1,141	15.0	1,234	18.3	1,330
Wealth quintile								
Lowest	26.3	2,146	27.1	2,111	26.6	2,182	24.4	1,771
Second	23.2	1,894	22.9	1,786	21.5	1,762	20.0	1,453
Middle	19.8	1,618	17.7	1,381	18.2	1,492	18.8	1,362
Higher	18.2	1,489	16.1	1,253	17.3	1,418	17.3	1,252
Highest	12.6	1,027	16.2	1,259	16.4	1,346	19.5	1,415
Birth order								
1	19.0	1,555	27.5	2,140	33.9	2,779	38.6	2,802
2-3	36.3	2,964	39.3	3,063	42.9	3,518	45.2	3,277
4-5	21.8	1,778	19.4	1,512	14.8	1,215	11.7	846
6+	23.0	1,877	13.8	1,074	8.4	689	4.5	328
Total	100.0	8,175	100.0	7,789	100.0	8,200	100.0	7,253

¹ Age of child refers to current age and does not include deceased children. Other characteristics refer to child's status at birth.

3.2.1 Size at Birth


Infants born with low birth weight (less than 2,500 grams) are considered to have a higher risk for early childhood death compared with babies born at an average or higher weight. For this analysis, mother's estimate of baby's size at birth was used as a proxy for birth weight. Babies who were reported by their mothers as "very small" or "smaller than average" are considered to have low birth weight. This alternative way of examining infant health is beneficial in Cambodia due to the lack of weight measurements in the earliest survey examined; thus mother's estimation was the largest source of information for this indicator (Figure 11). According to the 2000 CDHS, 83% of infants were not weighed at birth, compared with only 9% in 2014. Research shows that mother's estimation of a smaller than average birth size is positively associated with stunting later in childhood (Assaf, Kothari, and Pullum 2015) and therefore is a good proxy indicator for health.


Figure 11. Percentage of children born in the last 5 years who were not weighed, Cambodia 2000, 2005, 2010, and 2014 DHS



Based on mother's estimates, the proportion of children who were smaller than average size at birth showed a slight but significant decrease of 2 percentage points from 2000 to 2014 (Figure 12). Nearly all of the decrease in low birth weight infants occurred between 2005 and 2010. Between these two surveys, infants in the Plateau region, in the lowest wealth quintiles, or born to mothers with no education had the largest significant decreases in reported small size at birth. There were fewer infants reported as small in urban areas compared with rural areas and among mothers who do not smoke cigarettes compared with mothers who do, though smoking was not a significant factor in 2010. Appendix Table 8 shows confidence intervals and p-values that correspond to Figure 12.

Figure 12. Percentage of live births in the last 5 years who were estimated to be smaller than average by mothers, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS






Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * <0.05 , ** <0.01 , *** <0.001 .


3.2.2 Early Initiation of Breastfeeding

Early initiation of breastfeeding is important for the health of both mother and child. The World Health Organization (WHO) recommends that the newborn begin breastfeeding in the first hour. When given in the first hour, breast milk ensures that an infant receives the colostrum, the nutritious and rich first milk from the breast that contains antibodies that protect the infant in early and later life. Additionally, early feeding can facilitate the contraction of the uterus and lessen postpartum bleeding for the mother (Berkat and Sutan 2014).

Overall, Cambodia has seen a significant increase in early initiation of breastfeeding (in the first hour). Coverage of this indicator rose from 11% in 2000 to 66% in 2010 (Figure 13, Appendix Table 9). In the last survey interval, 2010-2014, coverage remained high, at 63%. Among subgroups, there was little variation in the coverage of breastfeeding in the first hour in the 2000, 2005, and 2010 CDHS. In 2014, some subgroups showed significant disparities that were not present in earlier surveys. Gaps in early initiation of breastfeeding were found by place of birth, mother's education, and wealth quintile. By region, infants in Phnom Penh had the lowest percentage of any subgroup receiving breast milk in the first hour, at 37% in 2014.

Figure 13. Percentage of last-born children in the 2 years preceding the survey who started breastfeeding within 1 hour of birth, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS






Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * <0.05 , ** <0.01 , *** <0.001 . Children with don't know or missing information are included within the "other" category for place of delivery.

3.2.3 Full Immunization


WHO considers a child to be fully vaccinated as having received vaccinations against tuberculosis (BCG); three doses against diphtheria, pertussis, tetanus (DPT) and polio; and against measles by the age of 12 months. Information on coverage of these vaccines originates from the mother's report of vaccination or the record of vaccination(s) on a health card. The coverage of each vaccine has increased in every survey (Figure 14). The results presented here reflect the proportion of fully vaccinated children among children age 12-23 months.


Figure 14. Percentage of children age 12-23 months who received each vaccine, Cambodia 2000, 2005, 2010, and 2014 DHS



The coverage of children age 12-23 months and fully vaccinated nearly doubled over the decade, from 40% in 2000 to 79% in 2010 (Figure 15). No improvement occurred between the surveys in 2010 and 2014, however; coverage dropped to 73% in 2014. In 2000, the proportion of fully vaccinated children differed by region, wealth quintile, mother's education, and place of delivery. Though improvements occurred for every subgroup by 2014, the gaps within subgroups have persisted. Compared with other subgroups, percentages of vaccinated children are higher in Phnom Penh and the Great Lake region, in the two wealthiest quintiles, among children with educated mothers, and among children born in a health facility. Appendix Table 10 contains supplementary results.

Figure 15. Percentage of children age 12-23 months who are fully immunized, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS





Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * <0.05 , ** <0.01 , *** <0.001 . Children with don't know or missing information are included within the "other" category for place of delivery.

3.2.4 Care Seeking for Diarrhea


Diarrhea—the second leading cause of death for children under age 5—is considered easily treatable with ORT. Interviewed mothers provide information about care seeking and treatment for all children under age 5 who had diarrhea in the 2 weeks preceding the survey. We present the following indicators of care seeking (treatment seeking and ORT) among the subset of children who recently experienced diarrhea, which ranged between 13% and 20% for any given survey.

Among children who experienced diarrhea in the previous 2 weeks, we considered children to have sought care if they were taken to a public or private facility, visited a health provider, or had a home visit from a provider. Pharmacies, shops, and traditional healers are not included in this definition. Care seeking among children tripled over the decade, from 22% in 2000 to 59% in 2010, before slightly decreasing in 2014 (Figure 17). In 2000, place of residence, wealth, mother's education, assistance at delivery, and age were significantly associated with seeking care for diarrhea. All subgroups experienced an increase in care

seeking, and the only significant disparities that remained in 2014 were by place of residence and place of delivery. (See Appendix Table 11 for additional information.)

Figure 16. Among children under age 5 who experienced diarrhea in the 2 weeks before the survey, percentage for whom treatment was sought from a health facility or provider, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS







Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * <0.05 , ** <0.01 , *** <0.001 . Children with don't know or missing information are included within the "other" category for place of delivery.

3.2.5 Treatment for Diarrhea

ORT is considered the best treatment for childhood diarrhea. This therapy includes children who were given ORT or who received increased fluids given by their mothers when the child had diarrhea. In 2000, among children under age 5 who experienced diarrhea in the 2 weeks before the survey, 62% received some form of ORT (Figure 17, Appendix Table 12). Over the 14-year survey period, ORT coverage remained stable at 52%-58% of the affected group of children. Significant differences in coverage between subgroups were found only in the 2005 survey. In this survey year, region, wealth, place of delivery, and current age significantly affected a child's likelihood of having received ORT. More recent survey years show quite uniform coverage among different subgroups, though diarrhea treatment remains lower in the Plain, Coastal, and Plateau regions and lowest among infants under age 6 months.

Figure 17. Among children under age 5 who experienced diarrhea in the 2 weeks before the survey, percentage given ORT or increased fluids, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS






Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * <0.05 , ** <0.01 , *** <0.001 . Children with don't know or missing information are included within the "other" category for place of delivery.

3.2.6 Nutritional Indices

A child's nutritional status affects the risk of disease, impaired physical growth, mental development, and premature death. Based on height, weight, and age, three indices are used to describe a child's nutritional status. The height-for-age index, a useful marker of health, is not sensitive to recent nutritional insufficiencies but instead reflects cumulative growth deficiencies and linear growth retardation. Low height-for-age, known as stunting, reflects malnutrition over an extended period and can be affected by long-term or chronic morbidity. The weight-for-height index reveals malnourished children at the time of the survey and the weight-for-age index reflects underweight children. Figure 18 shows the trends in these three indicators; we only examine stunting by background characteristics, however. Overall, we see improvement in every nutritional index, with the largest gains between the 2000 and 2005 CDHS.


Figure 18. Among children under age 5, percentage who are stunted, wasted, or underweight, Cambodia 2000, 2005, 2010, and 2014 DHS









Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * <0.05 , ** <0.01 , *** <0.001 .

The percentage of children who are stunted in Cambodia significantly decreased over the period of the four surveys, from 50% in 2000 to 32% in 2014 (Figure 19). The characteristics of the subgroups examined show significant differences in the percentages of stunted children. Though all subgroups saw a decrease over the period, the gaps remained consistent by rural-urban residence, region, wealth, mother's education, and age. In 2014, children in urban areas, in the Phnom Penh region, in the wealthiest quintile, children whose mothers have a secondary education, and those under age 1 had the smallest percentages stunted (see Appendix Table 13).

Figure 19. Among children under age 5, percentage who are stunted, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS





Note: A solid line indicates a significant change between two surveys, while a dotted line indicates no significant change. Significant change between 2000 and 2014 is indicated in the legend with asterisks to represent the p-value: * <0.05 , ** <0.01 , *** <0.001 .

4 DISCUSSION

4.1 Maternal Health

Overall, the trends in maternal and child health in Cambodia are encouraging. The results show a significant improvement in all maternal and child health indicators from the first CDHS survey in 2000 to the most recent in 2014, despite some stagnation or modest decline between 2010 and 2014 in certain indicators and among some subgroups. Indicators of ANC showed substantial improvement, with an increase of 70-80 percentage points in coverage, mostly before 2010. Disparities in coverage between regions, wealth quintiles, and education levels persisted over the survey period. Among women who sought maternal care, the quality of care also improved; however, these improvements leveled off between 2010 and 2014. Disparities in one of the components of good maternal care—discussing signs of complications with a provider—diminished over time. This suggests that once women obtain care, providers are giving less unequal treatment by place of residence, region, and wealth.

In Cambodia, indicators of care at birth and immediately after birth have also improved, including facility delivery, SBA, C-section, and PNC, with delivery in a health facility demonstrating the most impressive improvements. C-section rates have increased, although they are rare, performed for only 3% of all births in 2014. When needed and properly performed, C-sections can help avert mortality and morbidities that would otherwise accompany complications during delivery. WHO suggests that a C-section may be medically indicated for 10% to 15% of births; however, the surgery itself poses additional maternal and perinatal risks, particularly in facilities with sub-standard readiness to perform the surgery safely (Betran et al. 2016). For both SBA and PNC, there are pronounced disparities by sociodemographic groups, although these diminished from the first survey in 2000 to the most recent in 2014.

The continuum of care, beginning with ANC, is an instrumental factor in improving later health outcomes for both mother and child. The education and support a mother receives during ANC visits are critical to identifying early warning signs and encouraging use of health services later in the reproductive life cycle. Use of ANC services can influence more proximate outcomes during pregnancy and birth as well as child health outcomes, such as nutritional status (Alexander and Korenbrot 1995). This report shows the importance of receiving ANC to the continuum of maternal health care services—from ANC in pregnancy to skilled assistance in delivery and check-ups in the postpartum period. Women in Cambodia who had four or more ANC visits were more likely to have an SBA at most recent birth. Women who had four or more ANC visits or who had an SBA at birth were also more likely to receive PNC for the most recent birth in the previous 2 years. The percentage of women with four or more ANC visits, SBA, and PNC—the full continuum of care—increased substantially over the survey period, while the percentage with none of those services fell to negligible proportions by 2014.

These findings highlight the effects of Cambodia's efforts toward improving maternal health. In the mid-2000's, the Government of Cambodia launched a number of initiatives aimed at increasing use of maternal health services. These included Health Equity Fund (HEF) schemes to waive fees for the poor with reimbursement paid to health facilities by the HEF, voucher schemes to cover the cost of fees for ANC, assisted delivery, and PNC services in public health facilities, and the Midwife Incentive Scheme, which incentivized SBAs (Flores et al. 2013; Van de Poel et al. 2014). The Midwife Incentive Scheme allocated payment to trained professionals for live births delivered in health facilities. These were largely joint

initiatives between the Ministry of Health and the Ministry of Economy and Finance (Cambodia (MOH) and WHO 2014), with sponsorship by both the government and multiple international donor organizations. These efforts have contributed to the increasing use of maternal health services (Ir et al. 2010).

4.2 Child Health

Child health in Cambodia has seen significant and dramatic improvements since 2000. Large and significant increases in coverage occurred in early initiation of breastfeeding, full immunization of children age 12-23 months, and care seeking for children under age 5 who experienced diarrhea in the 2 weeks preceding the survey. Smaller but significant improvements occurred in the percentage of children smaller than average at birth and in the percentage of children stunted.

Rural mothers were more likely to report their child as smaller than average at birth, though rural areas had a larger and more significant decrease in small birth size than urban areas over the four surveys. Urban dwellers had a significantly smaller proportion of children stunted. Rural residents had larger and more significant gains in other indicators over the four surveys, such as care seeking for diarrhea, for which rural residents were more likely than their urban counterparts to seek care and had twice the coverage increase from 2000 to 2014.

The disparities between the lowest and highest wealth quintiles are significant in many health indicators in any survey year and in changes in indicator coverage over time. Poorer children saw steady improvement or upheld levels of coverage over the four surveys. In 2014, they were more likely to breastfeed within 1 hour of birth compared with children in wealthy households, though less likely to have all immunizations at age 1 and more likely to exhibit stunting. Having a more educated mother is significantly associated only with a child being fully immunized and exhibiting no stunting. Characteristics of the delivery—whether a doctor, nurse, or midwife assisted or whether the delivery took place at a health facility—were significantly associated with higher coverage of early breastfeeding, full immunization, care seeking for diarrhea, and ORT.

Cambodia's positive gains in child health may reflect the health promotion campaigns between 2000 and 2014. National campaigns using mass media to promote breastfeeding have resulted in significant and noteworthy growth in the percentage of women who breastfeed in the first hour after giving birth (Prak et al. 2014). In 2010, coverage was three to five times the level in 2000, though regionally progress stagnated or significantly decreased between 2010 and 2014. Moreover, promotion of breast milk substitutes (BMS) is illegal, though another study of the CDHS has shown that the use of BMS has greatly increased in urban areas (Forsberg, Sreeramareddy, and Low 2017). This may help explain the disparity shown in Phnom Penh, which has the smallest percentage of women breastfeeding in the first hour (Forsberg, Sreeramareddy, and Low 2017).

Cambodia has also focused efforts on effective immunization, which reduces levels of child mortality and decreases the number of reportable infectious disease cases. Full immunization coverage in children age 12-23 months significantly increased from 2000 to 2010, before dropping somewhat in 2014. This decline in the most recent survey largely results from the decrease in coverage of the measles vaccination and the continuing series of vaccinations against polio (Series 2 and 3) and diphtheria-tetanus-pertussis (Series 3). Though Cambodia has had success through the Reaching Every District strategy to

increase child immunization, in 2014 rural children, poor children, and children in the Plain and Plateau regions experienced significant decreases in immunization coverage (Chan Soeung et al. 2012).

Stunting among Cambodian children measured in the CHDS has been the focus of several studies that have showed associations with poverty, birth interval, little maternal education, and maternal smoking (Ikeda, Irie, and Shibuya 2013; Zanello, Srinivasan, and Shankar 2016). Although Cambodia did not meet the MDG target for stunting by 2015, the 17% significant decrease achieved in stunting from 2000 to 2014 is promising for meeting other global targets, such as the World Health Assembly Resolution's to reduce stunting by 40% by 2025 (Zanello, Srinivasan, and Shankar 2016). This analysis shows that gains were made in every subgroup over the 14-year period. Though little progress occurred between 2005 and 2010, between 2010 and 2014 there were large and significant decreases for all subgroups, with the largest improvement in the Coastal region. These consistent gains reflect the impact of the several national programs that have focused on improving nutrition over the past decade; nevertheless, regional and wealth disparities persist and should be addressed.

4.3 Conclusion

Maternal, newborn, and child mortality is largely preventable. The Sustainable Development Goals (SDGs) aim to end these preventable deaths by 2030 through universal access to good-quality health care (UNGA 2015). To meet these goals, monitoring trends in coverage of key indicators of healthcare interventions is critical for policymakers and program managers to decide where to concentrate resources. This report shows significant improvement in Cambodia's maternal and child health indicators over the past 2 decades, as evidenced by CDHS data. These positive changes reflect Cambodia's focused efforts in improving women's access to critical health services. Signs of stagnation in some indicators in recent years, however, suggest a possible loss of momentum. In order to meet the SDG targets, revitalized efforts may be required.

APPENDIX

Appendix Table 1. National-level trends for maternal health indicators, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000			2005			2010			2014			Difference ¹		
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	%	95% CI	2005-2005	2005-2010	2010-2014
Four or more antenatal care visits (ANC)	8.9	[7.7,10.1]	27.0	[24.9,29.2]	59.4	[57.0,61.7]	75.6	[73.6,77.5]	18.1***		32.4***		16.2***		66.7***
Timing of first ANC (before 4 months)	10.0	[8.8,11.3]	23.1	[21.4,25.0]	59.4	[57.3,61.4]	79.0	[77.0,80.8]	13.1***		36.3***		19.6***		69.0***
Mother took iron syrup/tablets during pregnancy	20.7	[18.9,22.7]	62.8	[60.3,65.2]	89.4	[88.1,90.5]	95.6	[94.6,96.4]	42.1***		26.6***		6.2***		74.9***
Mother took deworming medication during pregnancy	n/a		10.7	[9.6,11.8]	44.5	[42.5,46.5]	72.2	[70.3,73.9]	n/a		33.8***		27.7***		61.5***
Blood pressure checked during ANC	61.3	[58.5,64.0]	80.6	[78.8,82.3]	90.6	[89.4,91.6]	96.1	[95.2,96.8]	19.3***		10.0***		5.5***		34.8***
Informed of pregnancy complications during ANC	40.1	[37.6,42.7]	60.3	[58.2,62.4]	80.0	[78.3,81.6]	82.1	[80.1,84.0]	20.2***		19.7***		2.1		42.0***
Birth delivered in a facility	9.9	[8.8,11.2]	21.5	[19.6,23.5]	53.8	[51.1,56.4]	83.2	[80.9,85.2]	11.6***		32.3***		29.4***		73.3***
Births assisted by a skilled birth attendant (SBA)	31.8	[29.0,34.8]	43.8	[41.1,46.5]	71.0	[68.6,73.4]	89.0	[87.1,90.7]	12.0***		27.2***		18.0***		57.2***
Births delivered by Caesarean section	0.8	[0.6,1.2]	1.8	[1.5,2.3]	3.0	[2.5,3.6]	6.3	[5.6,7.1]	1.0***		1.2***		3.3***		5.5***
Postnatal care for the mother	n/a		64.7	[62.2,67.1]	74.2	[71.9,76.4]	90.4	[88.5,92.0]	n/a		9.5***		16.2***		25.7***

¹ Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions, p-values *<0.05, **<0.01, ***<0.001.

Appendix Table 2. Percentage of women with four or more antenatal care visits for their most recent pregnancy among women age 15–49 with a live birth in the 5 years preceding the survey, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000			2005			2010			2014			Difference ¹			
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014
Total	8.9	[7.7,10.1]		27.0	[24.9,29.2]		59.4	[57.0,61.7]		75.6	[73.6,77.5]		18.1***	32.4***	16.2***	66.7***
Place of residence																
Urban	23.0	[19.3,27.1]	***	43.2	[38.6,48.0]	***	80.3	[77.4,82.9]	***	85.4	[82.8,87.7]	***	20.2***	37.1***	5.1**	62.4***
Rural	6.6	[5.5,8.0]		24.4	[22.1,26.8]		55.3	[52.6,58.0]		73.9	[71.6,76.1]		17.8***	30.9***	18.6***	67.3***
Region																
Phnom Penh	38.8	[32.1,46.1]	***	54.2	[44.3,63.8]	***	85.8	[81.7,89.1]	***	88.0	[83.7,91.2]	***	15.4*	31.6***	2.2	49.2***
Plain	6.1	[4.4,8.4]		24.8	[21.1,29.0]		57.7	[53.3,62.0]		77.2	[73.4,80.6]		18.7***	32.9***	19.5***	71.1***
Great Lake	10.2	[8.4,12.5]		29.8	[26.5,33.4]		62.6	[58.6,66.5]		80.2	[76.9,83.1]		19.6***	32.8***	17.6***	70.0***
Coastal	2.6	[1.2,5.5]		19.2	[15.6,23.3]		46.5	[40.1,53.1]		71.5	[65.8,76.5]		16.6***	27.3***	25.0***	68.9***
Plateau	4.5	[3.0,6.7]		15.1	[12.7,18.0]		48.1	[43.0,53.2]		60.2	[55.4,64.9]		10.6***	33.0***	12.1***	55.7***
Education																
None	3.7	[2.7,5.1]	***	15.6	[13.3,18.3]	***	40.0	[36.0,44.1]	***	54.9	[49.5,60.2]	***	11.9***	24.4***	14.9***	51.2***
Primary	7.2	[6.0,8.8]		24.6	[22.6,26.9]		56.8	[53.9,59.7]		74.0	[71.7,76.1]		17.4***	32.2***	17.2***	66.8***
Secondary+	26.5	[22.8,30.5]		50.1	[45.4,54.7]		77.7	[74.9,80.2]		86.1	[84.3,87.8]		23.6***	27.6***	8.4***	59.6***
Wealth quintile																
Lowest	2.9	[2.0,4.3]	***	15.1	[12.6,17.9]	***	42.8	[38.9,46.9]	***	60.5	[55.8,65.0]	***	12.2***	27.7***	17.7***	57.6***
Second	5.5	[4.1,7.4]		19.0	[16.5,21.8]		51.4	[47.4,55.5]		69.6	[65.5,73.4]		13.5***	32.4***	18.2***	64.1***
Middle	4.8	[3.4,6.6]		22.5	[19.4,25.8]		57.9	[54.0,61.8]		76.6	[73.0,79.9]		17.7***	35.4***	18.7***	71.8***
Fourth	8.1	[5.9,10.9]		30.6	[26.5,35.1]		70.6	[66.3,74.6]		85.8	[83.2,88.0]		22.5***	40.0***	15.2***	77.7***
Highest	31.3	[27.4,35.5]		56.8	[52.4,61.1]		82.5	[79.7,84.9]		88.9	[86.7,90.8]		25.5***	25.7***	6.4***	57.6***
Mother's age at birth																
<20	10.9	[7.9,15.0]	**	28.4	[23.8,33.4]	***	62.9	[57.8,67.8]	***	72.1	[67.4,76.4]	***	17.5***	34.5***	9.2*	61.2***
20-34	9.5	[8.3,10.9]		29.4	[27.1,31.8]		62.2	[59.8,64.6]		77.6	[75.6,79.5]		19.9***	32.8***	15.4***	68.1***
35-49	6.2	[4.7,8.1]		18.3	[15.7,21.2]		43.4	[38.9,48.0]		63.7	[58.4,68.7]		12.1***	25.1***	20.3***	57.5***
Birth order																
1	13.0	[10.5,15.9]	***	39.4	[35.8,43.2]	***	71.9	[69.0,74.6]	***	81.3	[78.9,83.5]	***	26.4***	32.5***	9.4***	68.3***
2-3	11.1	[9.4,13.1]		29.2	[26.5,32.1]		60.9	[58.2,63.5]		78.6	[76.4,80.6]		18.1***	31.7***	17.7***	67.5***
4-5	6.6	[4.8,8.9]		17.4	[14.8,20.2]		47.1	[42.3,51.9]		63.4	[58.0,68.4]		10.8***	29.7***	16.3***	56.8***
6+	4.7	[3.3,6.8]		13.7	[10.9,17.1]		29.0	[24.2,34.2]		35.4	[28.7,42.7]		9.0***	15.3***	6.4	30.7***
Total	5,714			5,865			6,472			5,973						

¹ Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions, p-values * <0.05 , ** <0.01 , *** <0.001 .

Appendix Table 3. Among women age 15-49 with a live birth in the 5 years preceding the survey who received antenatal care for their most recent birth, percentage who were informed of signs of complications, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000			2005			2010			2014			Difference ¹			
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014
Total	40.1	[37.6,42.7]		60.3	[58.2,62.4]		80.0	[78.3,81.6]		82.1	[80.1,84.0]		20.2***	19.7***	2.1	42.0***
Place of residence																
Urban	47.9	[41.7,54.2]	**	64.2	[60.0,68.2]		87.9	[85.6,89.9]	***	83.1	[80.5,85.4]		16.3***	23.7***	-4.8**	35.2***
Rural	38.1	[35.3,41.0]		59.6	[57.2,61.9]		78.3	[76.3,80.2]		82.0	[79.6,84.1]		21.5***	18.7***	3.7*	43.9***
Region																
Phnom Penh	60.9	[51.3,69.8]	***	68.6	[60.1,75.9]	***	92.0	[87.8,94.8]	***	84.6	[80.4,88.0]		7.7	23.4***	-7.4**	23.7***
Plain	41.0	[36.2,45.9]		55.5	[51.9,59.0]		75.2	[71.7,78.4]		80.9	[76.2,84.9]		14.5***	19.7***	5.7	39.9***
Great Lake	33.2	[29.7,36.9]		67.2	[63.9,70.4]		84.4	[82.0,86.5]		84.7	[82.3,86.8]		34.0***	17.2***	0.3	51.5***
Coastal	50.2	[40.3,60.1]		60.6	[54.2,66.7]		75.7	[68.6,81.7]		82.4	[78.1,86.1]		10.4	15.1**	6.7	32.2***
Plateau	34.0	[28.2,40.4]		49.3	[43.8,54.7]		78.4	[75.0,81.5]		78.6	[74.9,81.9]		15.3***	29.1***	0.2	44.6***
Wealth quintile																
Lowest	31.5	[26.8,36.6]	***	55.2	[50.4,59.9]	**	74.6	[70.8,78.1]	***	78.7	[73.2,83.3]		23.7***	19.4***	4.1	47.2***
Second	32.8	[28.1,37.8]		57.3	[53.2,61.4]		77.4	[73.7,80.7]		81.1	[77.7,84.1]		24.5***	20.1***	3.7	48.3***
Middle	36.3	[30.7,42.4]		62.3	[57.8,66.5]		81.5	[78.1,84.4]		83.2	[79.8,86.1]		26.0***	19.2***	1.7	46.9***
Fourth	41.7	[36.1,47.6]		59.5	[55.4,63.4]		81.7	[78.3,84.6]		85.0	[82.0,87.7]		17.8***	22.2***	3.3	43.3***
Highest	52.6	[47.1,57.9]		67.3	[62.9,71.4]		85.9	[82.7,88.5]		83.1	[79.5,86.2]		14.7***	18.6***	-2.8	30.5***
Education																
None	27.6	[23.9,31.6]	***	56.5	[51.3,61.5]	**	74.4	[69.8,78.5]	***	76.7	[71.3,81.4]	**	28.9***	17.9***	2.3	49.1***
Primary	40.3	[37.0,43.7]		58.9	[56.4,61.3]		78.8	[76.6,80.8]		81.7	[79.3,83.9]		18.6***	19.9***	2.9	41.4***
Secondary+	52.9	[47.7,58.0]		67.1	[62.8,71.2]		85.4	[83.0,87.6]		84.5	[82.0,86.7]		14.2***	18.3***	-0.9	31.6***
Mother's age at birth																
<20	42.1	[34.6,50.0]		57.7	[51.9,63.3]	*	74.2	[69.2,78.6]	**	76.4	[70.5,81.5]	*	15.6**	16.5***	2.2	34.3***
20-34	40.9	[37.9,43.9]		59.2	[56.8,61.6]		80.3	[78.4,82.1]		82.7	[80.8,84.5]		18.3***	21.1***	2.4	41.8***
35-49	36.8	[31.6,42.3]		66.0	[61.5,70.2]		82.0	[78.6,85.0]		83.6	[78.3,87.8]		29.2***	16.0***	1.6	46.8***
Birth order																
1	42.5	[37.3,48.0]		59.9	[56.3,63.4]		79.2	[76.4,81.7]		80.6	[77.4,83.4]		17.4***	19.3***	1.4	38.1***
2-3	40.5	[36.4,44.6]		58.3	[55.0,61.4]		80.7	[78.4,82.8]		84.2	[82.1,86.1]		17.8***	22.4***	3.5*	43.7***
4-5	41.5	[36.8,46.4]		64.2	[60.0,68.3]		80.3	[76.2,83.8]		80.3	[75.4,84.4]		22.7***	16.1***	0.0	38.8***
6+	35.1	[30.3,40.4]		62.7	[57.1,67.9]		79.0	[73.5,83.6]		75.7	[65.3,83.8]		27.6***	16.3***	-3.3	40.6***
Total	2,543			4,213			5,804			5,704						

¹ Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions, p-values * <0.05 , ** <0.01 , *** <0.001 .

Appendix Table 4. Percentage of births that were assisted by a skilled birth attendant, among all births in the 5 years preceding the survey to women age 15-49, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000			2005			2010			2014			Difference ¹			
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014
Total	31.8	[29.0,34.8]		43.8	[41.1,46.5]		71.0	[68.6,73.4]		89.0	[87.1,90.7]		12.0***	27.2***	18.0***	57.2***
Place of residence																
Urban	57.2	[49.7,64.4]	***	70.1	[64.2,75.4]	***	94.7	[92.9,96.1]	***	97.8	[96.5,98.6]	***	12.9**	24.6***	3.1**	40.6***
Rural	28.0	[25.0,31.2]		39.5	[36.5,42.5]		66.6	[63.8,69.3]		87.6	[85.4,89.5]		11.5***	27.1***	21.0***	59.6***
Region																
Phnom Penh	88.9	[81.7,93.5]	***	86.0	[73.2,93.2]	***	98.8	[97.3,99.4]	***	96.1	[92.3,98.1]	***	-2.9	12.8***	-2.7*	7.2*
Plain	32.4	[27.1,38.2]		49.7	[44.6,54.8]		75.7	[72.0,79.2]		94.6	[91.7,96.5]		17.3***	26.0***	18.9***	62.2***
Great Lake	25.2	[21.4,29.5]		36.5	[32.0,41.3]		67.4	[62.5,72.0]		91.2	[88.8,93.1]		11.3***	30.9***	23.8***	66.0***
Coastal	30.7	[22.6,40.3]		46.4	[39.2,53.9]		70.7	[62.2,78.1]		93.1	[87.7,96.3]		15.7*	24.3***	22.4***	62.4***
Plateau	24.1	[19.7,29.2]		20.0	[16.2,24.3]		51.7	[45.1,58.1]		69.6	[62.7,75.7]		-4.1	31.7***	17.9	45.5***
Wealth quintile																
Lowest	14.7	[11.9,17.9]	***	20.7	[17.7,24.1]	***	48.7	[44.6,52.8]	***	75.2	[69.9,79.8]	***	6.0**	28.0***	26.5***	60.5***
Second	21.3	[17.8,25.2]		29.0	[25.5,32.8]		63.7	[59.8,67.4]		87.0	[84.1,89.5]		7.7**	34.7***	23.3***	65.7***
Middle	27.4	[23.5,31.7]		39.6	[35.9,43.5]		74.5	[70.5,78.1]		92.7	[90.2,94.6]		12.2***	34.9***	18.2***	65.3***
Fourth	40.7	[36.2,45.4]		61.9	[57.6,66.1]		86.5	[83.6,89.0]		96.5	[94.6,97.7]		21.2***	24.6***	10***	55.8***
Highest	81.2	[76.5,85.1]		89.9	[86.4,92.6]		96.7	[94.9,97.9]		98.4	[97.1,99.1]		8.7***	6.8***	1.7	17.2**
Mother's education																
None	19.3	[16.3,22.7]	***	22.0	[18.9,25.5]	***	46.9	[42.5,51.4]	***	71.8	[65.5,77.3]	***	2.7	24.9***	24.9***	52.5***
Primary	30.9	[27.8,34.2]		42.5	[39.7,45.4]		70.4	[67.6,73.0]		88.5	[86.5,90.3]		11.6***	27.9***	18.1***	57.6***
Secondary+	65.8	[61.0,70.4]		79.5	[76.0,82.6]		90.5	[88.6,92.2]		97.0	[96.0,97.8]		13.7***	11.0***	6.5***	31.2***
Mother's age at birth																
<20	34.5	[29.1,40.2]	**	44.2	[39.4,49.1]	*	73.1	[68.7,77.2]	***	89.3	[85.9,92.0]	***	9.7**	28.9***	16.2***	54.8***
20-34	33.1	[30.1,36.2]		44.9	[41.9,47.9]		72.2	[69.7,74.6]		89.9	[88.0,91.5]		11.8***	27.3***	17.7***	56.8***
35-49	26.2	[22.6,30.2]		39.3	[35.6,43.0]		63.0	[58.7,67.1]		81.3	[76.8,85.1]		13.1***	23.7***	18.3***	55.1***
Birth order																
1	43.4	[39.5,47.4]	***	54.2	[50.8,57.5]	***	80.4	[78.0,82.6]	***	94.1	[92.6,95.3]	***	10.8***	26.2***	13.7***	50.7***
2-3	36.0	[32.5,39.7]		45.8	[42.5,49.3]		72.1	[69.2,74.8]		89.4	[87.6,91.0]		9.8***	26.3***	17.3***	53.4***
4-5	26.7	[23.4,30.3]		34.7	[31.3,38.3]		58.7	[54.1,63.1]		80.1	[75.1,84.2]		8.0**	24.0***	21.4***	53.4***
6+	20.4	[17.3,24.0]		29.9	[25.8,34.3]		49.6	[44.1,55.0]		65.0	[56.8,72.4]		9.5***	19.7***	15.4**	44.6***
Number of ANC visits^{2,3}																
0-3 visits	30.8	[27.9,33.7]	***	37.9	[35.0,40.8]	***	55.8	[52.3,59.2]	***	74.7	[70.1,78.8]	***	7.1***	17.9***	18.9***	43.9***
4 or more visits	72.3	[65.9,78.0]		69.6	[66.0,73.0]		86.4	[84.4,88.1]		95.5	[94.2,96.4]		-2.7	16.8***	9.1	23.2***
Place of delivery³																
Home, other	24.4	[21.8,27.2]	***	28.4	[25.7,31.3]	***	37.8	[34.4,41.3]	***	35.4	[29.0,42.3]	***	4.0*	9.4***	-2.4	11.0**
Health facility	99.2	[97.9,99.7]		99.7	[99.2,99.9]		99.6	[99.2,99.8]		99.9	[99.8,100.0]		0.5	-0.1	0.3*	0.7**
Total	8,715			7,789			8,200			7,253						

¹ Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions, p-values <0.05. **<0.01, ***<0.001. ² Number of ANC visits is calculated among most recent births only. ³ Births with don't know or missing information are included with the category "0-3 visits" for ANC and the "home, other" category for place of delivery.

Appendix Table 5. Percentage of women who received a postnatal check-up within 2 days of delivering their most recent child among women age 15-49 with a live birth in the 2 years preceding the survey, according to background characteristics, Cambodia 2005, 2010, and 2014 DHS

Background characteristic	2005			2010			2014			Difference ¹		
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2005-2010	2010-2014	2000-2014
Total	64.7	[62.2,67.1]		74.2	[71.9,76.4]		90.4	[88.5,92.0]		9.5***	16.2***	25.7***
Place of residence			**			***			***			
Urban	73.5	[67.5,78.8]		88.4	[84.7,91.3]		98.1	[97.0,98.8]		14.9***	9.7***	24.6***
Rural	63.2	[60.5,65.9]		71.5	[68.8,74.0]		89.2	[87.0,91.0]		8.3***	17.7***	26.0***
Region			***			***			***			
Phnom Penh	84.6	[73.1,91.8]		95.1	[90.0,97.6]		100.0			10.5*	4.9***	15.4***
Plain	66.2	[61.7,70.5]		75.0	[70.6,78.9]		89.0	[84.6,92.2]		8.8**	14.0***	22.8***
Great Lake	65.0	[60.5,69.2]		77.0	[73.1,80.5]		94.4	[92.1,96.1]		12.0***	17.4***	29.4***
Coastal	55.8	[50.2,61.4]		77.8	[70.2,83.9]		94.3	[89.8,96.8]		22.0***	16.5***	38.5***
Plateau	53.1	[47.5,58.7]		53.3	[47.8,58.7]		80.0	[75.2,84.1]		0.2	26.7***	26.9***
Wealth quintile			***			***			***			
Lowest	53.9	[49.1,58.6]		60.9	[56.2,65.3]		84.3	[79.9,88.0]		7.0*	23.4***	30.4***
Second	57.8	[53.0,62.4]		66.3	[61.6,70.8]		87.7	[84.4,90.4]		8.5*	21.4***	29.9***
Middle	64.9	[60.3,69.3]		77.6	[72.6,81.9]		93.7	[90.4,96.0]		12.7***	16.1***	28.8***
Fourth	69.8	[64.4,74.7]		82.9	[78.4,86.7]		92.2	[87.9,95.1]		13.1***	9.3**	22.4***
Highest	85.7	[81.2,89.2]		91.7	[88.2,94.2]		95.6	[90.2,98.1]		6.0*	3.9	9.9***
Education			***			***			***			
None	56.5	[51.5,61.3]		56.6	[50.8,62.2]		79.8	[73.7,84.8]		0.1	23.2***	23.3***
Primary	62.2	[59.2,65.0]		73.6	[70.6,76.4]		90.3	[87.7,92.3]		11.4***	16.7***	28.1***
Secondary+	83.9	[80.0,87.1]		86.9	[83.8,89.5]		94.2	[92.1,95.7]		3.0	7.3***	10.3***
Mother's age at birth						**						
<20	61.1	[54.3,67.6]		78.3	[72.7,83.1]		88.7	[83.9,92.2]		17.2***	10.4**	27.6***
20-34	65.7	[62.8,68.5]		74.9	[72.5,77.2]		91.0	[88.9,92.7]		9.2***	16.1***	25.3***
35-49	62.5	[57.5,67.2]		65.3	[58.7,71.4]		87.1	[80.6,91.7]		2.8	21.8***	24.6***
Birth order			**			***			*			
1	69.5	[65.7,73.1]		80.5	[77.6,83.1]		91.9	[89.7,93.6]		11.0***	11.4***	22.4***
2-3	64.8	[61.4,68.0]		74.3	[71.0,77.4]		90.2	[87.5,92.4]		9.5***	15.9***	25.4***
4-5	60.8	[55.9,65.4]		68.5	[62.4,74.0]		88.2	[83.5,91.7]		7.7*	19.7***	27.4***
6+	59.2	[52.8,65.4]		49.3	[40.9,57.8]		82.8	[71.2,90.3]		-9.9	33.5***	23.6***
Number of ANC visits²			***			***			***			
0-3 visits	59.2	[56.3,62.0]		56.4	[52.7,60.1]		81.9	[77.8,85.4]		-2.8	25.5***	22.7***
4 or more visits	78.5	[75.2,81.5]		84.0	[81.8,86.1]		93.1	[91.1,94.7]		5.5**	9.1***	14.6***
Place of delivery²			***			***			***			
Home, other	56.8	[53.9,59.7]		41.9	[38.0,45.9]		57.1	[49.4,64.5]		-14.9***	15.2***	0.3
Health facility	87.8	[84.6,90.4]		90.6	[89.1,92.0]		94.6	[93.0,95.9]		2.8	4.0***	6.8***
Assistance at delivery²			***			***			***			
No one, other	48.2	[44.4,52.0]		33.2	[28.2,38.5]		51.1	[41.6,60.4]		-15.0***	17.9***	2.9
Doctor, nurse, midwife	82.0	[79.3,84.5]		84.9	[82.9,86.8]		93.5	[91.7,95.0]		2.9	8.6***	11.5***
Total	3,083			3,187			2,944					

¹ Percentage point difference between two surveys and between 2005 and 2014 with significant tests for the difference in proportions, p-values *<0.05, **<0.01, ***<0.001. ² Women with don't know or missing information are included with the category "0-3 visits" for ANC and the "home, other" category for place of delivery and the "no one, other" category for assistance at delivery.

Appendix Table 6. Distribution of women with antenatal care (four or more visits), a skilled birth attendant at delivery, or postnatal check-up within 2 days of delivering their most recent child among women age 15-49 with a live birth in the 2 years preceding the survey, according to background characteristics, Cambodia 2005, 2010, and 2014 DHS

	2005		2010		2014		Difference ¹		
	%	95% CI	%	95% CI	%	95% CI	2005-2010	2010-2014	2000-2014
No ANC4, no SBA, no PNC	22.7	[20.6,25.0]	10.5	[9.1,12.1]	2.4	[1.8,3.2]	-12.2***	-8.1***	-20.3***
Yes ANC4, no SBA, no PNC	3.8	[3.1,4.7]	3.4	[2.7,4.3]	1.2	[0.7,2.0]	-0.4	-2.2***	-2.6***
Yes ANC4, yes SBA, no PNC	2.3	[1.7,3.1]	6.9	[5.8,8.1]	4.0	[2.9,5.5]	4.6***	-2.9**	1.7*
Yes ANC4, yes SBA, yes PNC	17.9	[15.7,20.2]	51.3	[48.4,54.1]	69.4	[66.4,72.2]	33.4***	18.1***	51.5***
Yes ANC4, no SBA, yes PNC	4.5	[3.6,5.6]	2.7	[2.0,3.7]	1.2	[0.7,1.9]	-1.8**	-1.5**	-3.3***
No ANC4, yes SBA, no PNC	6.5	[5.4,7.7]	5.1	[4.1,6.3]	2.0	[1.4,3.0]	-1.4	-3.1***	-4.5***
No ANC4, no SBA, yes PNC	20.2	[17.7,22.8]	4.1	[3.2,5.4]	2.6	[1.7,3.9]	-16.1***	-1.5	-17.6***
No ANC4, yes SBA, yes PNC	22.2	[20.0,24.6]	16.0	[14.4,17.8]	17.3	[15.5,19.3]	-6.2***	1.3	-4.9**
Total	3,083		3,187		2,944				

¹ Percentage point difference between two surveys and between 2005 and 2014 with significant tests for the difference in proportions, p-values *<0.05, **<0.01, ***<0.001.

Appendix Table 7. National-level trends for child health indicators, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000		2005		2010		2014		Difference ¹				
	%	95% CI	%	95% CI	%	95% CI	%	95% CI	2005-2000	2010-2000	2005-2010	2010-2014	2000-2014
Mother's estimate of child's size at birth	13.8	[12.8,14.9]	14.4	[13.5,15.3]	11.4	[10.3,12.5]	11.6	[10.4,12.9]	0.6		-3.0*	0.2	-2.2*
Early initiation of breastfeeding in the first hour	11.1	[9.9,12.4]	35.5	[33.3, 37.9]	65.8	[63.3, 68.2]	62.6	[60.0, 65.0]	24.4*		30.3*	-3.2*	51.5*
Full immunization	39.9	[35.9,44.0]	66.6	[63.2,69.8]	78.9	[75.7,81.7]	73.4	[70.2,76.5]	26.7*		12.3*	-5.5*	33.5*
Care seeking for diarrhea	21.6	[18.8,24.7]	41.1	[37.5,44.8]	58.9	[55.2,62.6]	55.5	[50.8,60.0]	19.5*		17.8*	-3.4*	33.9*
Oral Rehydration Therapy (ORT) for diarrhea	74.1	[70.9,76.9]	58.4	[54.9,61.9]	52.6	[48.6,56.6]	57.0	[51.7,62.2]	-15.7*		-5.8*	4.4	-17.1*
Wasting	16.8	[15.0,18.7]	8.5	[7.5,9.7]	11.3	[10.1,12.7]	9.7	[8.6,10.9]	-8.2***		2.8**	-1.6	-7.0***
Underweight	38.4	[36.3,40.6]	28.2	[26.3,30.2]	27.8	[25.8,29.9]	24.1	[22.3,26.0]	-10.2***		0.4	-3.7*	-14.3***
Stunting	49.6	[47.3,51.9]	42.4	[40.0,44.7]	39.1	[36.9,41.4]	31.9	[30.0,33.9]	-7.2***		-3.3	-7.2***	-17.7***

¹ Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions, p-values *<0.05, **<0.01, ***<0.001.

Appendix Table 8. Percentage of children born in the last 5 years who were smaller than average by mother's estimation, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000			2005			2010			2014			Difference ¹			
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014
Total	13.8	[12.8,14.9]		14.4	[13.5,15.3]		11.4	[10.3,12.5]		11.6	[10.4,12.9]		0.6	-3.0***	0.2	-2.2**
Place of residence																
Urban	9.1	[6.8,12.0]	**	11.7	[10.0,13.6]	**	8.1	[6.5,10.2]	**	8.5	[6.6,10.7]	**	-2.6	3.6**	0.3	-0.6
Rural	14.5	[13.5,15.6]		14.9	[13.9,15.9]		12.0	[10.8,13.3]		12.1	[10.7,13.6]		0.3	-2.9***	0.1	-2.4**
Region																
Phnom Penh	6.6	[3.4,12.6]	**	5.9	[4.3,8.0]	***	5.5	[3.1,9.5]	**	7.1	[4.3,11.5]	***	-0.7	-0.4	1.6	0.5
Plain	13.9	[11.9,16.0]		11.1	[9.4,13.0]		11.3	[9.4,13.6]		9.9	[7.8,12.5]		-2.8*	0.3	1.4	4.0*
Great Lake	14.1	[12.7,15.6]		17.0	[15.6,18.6]		13.3	[11.5,15.4]		11.2	[9.2,13.6]		2.9**	-3.7**	-2.1	-2.9*
Coastal	10.9	[7.8,14.9]		13.1	[10.7,16.0]		9.7	[7.3,12.7]		10.9	[8.6,13.9]		2.2	3.4	-1.3	0.1
Plateau	17.6	[15.1,20.4]		22.7	[20.6,25.0]		11.4	[9.5,13.7]		17.8	[15.3,20.7]		5.1**	-11.3***	6.4***	0.2
Wealth quintile																
Lowest	14.4	[12.5,16.6]		18.6	[16.6,20.7]	***	13.4	[11.3,15.9]	*	13.9	[11.1,17.2]	*	4.1**	-5.1***	0.5	-0.5
Second	14.4	[12.1,17.0]		16.3	[14.1,18.8]		12.6	[10.4,15.1]		11.5	[9.4,14.0]		2.0	-3.8*	-1.1	-2.9
Middle	14.1	[11.9,16.6]		13.7	[11.6,16.1]		11.1	[8.8,14.0]		12.6	[9.9,15.9]		-0.4	-2.6	1.5	-1.5
Fourth	14.4	[11.8,17.3]		11.6	[9.3,14.3]		9.8	[7.9,12.1]		10.7	[8.6,13.3]		-2.8	1.8	-0.9	-3.7*
Highest	10.1	[7.7,13.2]		8.3	[6.5,10.7]		8.5	[6.4,11.2]		8.4	[6.7,10.5]		1.8	0.2	-0.1	1.7
Mother's education																
No education	14.8	[13.1,16.7]	*	18.7	[15.7,22.1]	***	13.6	[10.9,16.9]		15.6	[12.5,19.3]	*	3.9*	-5.0*	1.9	0.8
Primary	14.1	[12.6,15.6]		14.4	[13.1,15.7]		11.4	[10.2,12.8]		10.9	[9.4,12.8]		0.3	-3.0**	-0.5	-3.1**
Secondary+	10.4	[8.3,12.8]		8.5	[6.6,10.8]		9.7	[7.9,12.0]		10.8	[9.2,12.7]		1.9	1.3	-1.1	0.5
Mother's age at birth																
<20	11.8	[9.0,15.3]	**	16.4	[13.5,19.9]		12.7	[10.0,16.0]		13.8	[11.0,17.3]		4.6*	-3.8	1.1	2.0
20-34	13.2	[12.0,14.4]		13.9	[12.9,15.1]		11.1	[9.9,12.4]		11.1	[9.9,12.4]		0.8	-2.9**	0.0	-2.0*
35-49	16.9	[14.6,19.4]		15.0	[12.8,17.6]		12.2	[9.9,14.9]		12.6	[9.7,16.4]		-1.9	-2.9	0.5	-4.3
Mother smokes cigarettes																
No	13.5	[12.5,14.6]	*	13.9	[13.0,14.9]	***	11.2	[10.2,12.4]		11.2	[10.1,12.4]	***	0.4	-2.6***	0.0	-2.3**
Yes	17.9	[13.8,22.8]		24.3	[18.5,31.3]	***	16.5	[10.5,25.0]		23.2	[13.3,37.3]	***	6.4	-7.8	6.7	5.3
Birth order																
1	14.5	[12.5,16.8]		15.8	[14.1,17.7]	***	13.3	[11.6,15.2]	***	14.1	[12.2,16.2]	***	1.2	-2.5	0.8	-0.5
2-3	13.0	[11.5,14.8]		12.1	[10.8,13.6]		9.7	[8.4,11.1]		9.2	[7.9,10.8]		-0.9	2.4*	-0.5	3.8***
4-5	12.3	[10.6,14.1]		13.7	[11.6,16.1]		9.8	[7.8,12.1]		10.3	[8.1,13.1]		1.4	3.9*	-0.6	-1.9
6+	15.9	[13.8,18.1]		19.2	[16.4,22.4]		15.5	[11.7,20.2]		16.4	[11.8,22.2]		3.4*	-3.7	0.9	0.5
Birth interval																
First Birth	14.7	[12.6,17.0]		15.9	[14.1,17.8]	**	13.4	[11.7,15.3]	*	14.4	[12.5,16.5]	***	1.2	-2.5	1.0	-0.3
<24	14.4	[12.3,16.8]		16.2	[13.6,19.2]		10.3	[8.1,13.0]		13.1	[9.8,17.4]		1.8	-5.9**	2.8	-1.3
24-47	13.0	[11.6,14.5]		14.3	[12.8,15.9]		10.4	[8.8,12.1]		9.9	[8.2,11.9]		1.3	-3.9**	0.5	3.1*
48+	14.2	[12.2,16.5]		11.9	[10.2,13.8]		10.4	[8.8,12.2]		8.6	[7.1,10.3]		-2.3	-1.5	1.8	5.6***
Place of delivery²																
Home, other	13.9	[12.8,15.0]		15.9	[14.8,17.0]	***	13.4	[11.7,15.2]	***	17.3	[14.0,21.1]	***	2.0*	-2.5*	3.9*	3.4*
Health facility	13.0	[9.8,16.9]		9.1	[7.5,11.2]		9.7	[8.5,11.0]		10.4	[9.2,11.7]		3.8*	0.6	-0.7	-2.6

Continued...

Appendix Table 8—Continued

	2000			2005			2010			2014			Difference ¹			
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014
Assistance at delivery²																
No one, other	14.1	[12.9,15.4]		17.5	[16.1,19.1]		14.6	[12.5,17.0]		19.5	[15.2,24.6]		3.4***	-3.0*	4.9*	5.4*
Doctor, nurse, midwife	13.1	[11.5,15.0]		10.4	[9.2,11.8]		10.1	[9.0,11.3]		10.6	[9.4,11.9]		-2.8**	-0.3	0.5	-2.6*
Child's current age																
<6 mo	14.2	[11.6,17.2]		17.1	[14.4,20.1]		12.8	[10.0,16.2]		11.1	[8.5,14.4]		2.9	-4.3	-1.7	-3.0
6-11 mo	14.6	[11.8,17.9]		13.5	[10.6,17.0]		8.4	[6.3,11.2]		8.7	[6.6,11.4]		-1.1	5.1**	0.2	5.9**
1 years	14.9	[12.7,17.4]		14.5	[12.4,16.9]		10.4	[8.7,12.4]		11.5	[9.4,14.0]		-0.4	-4.1**	1.1	-3.4*
2 years	13.1	[11.1,15.2]		12.8	[10.8,15.1]		10.9	[8.6,13.6]		10.9	[8.9,13.2]		-0.3	-1.9	0.0	-2.2
3 years	12.2	[10.5,14.2]		13.3	[11.0,15.9]		12.0	[10.1,14.1]		9.7	[7.9,12.0]		1.1	-1.3	2.2	2.5
4 years	10.9	[9.3,12.8]		13.7	[11.8,15.9]		10.6	[8.6,13.0]		12.7	[10.5,15.2]		2.8*	-3.1*	2.0	1.7
Child's sex			**			*										
Male	12.5	[11.2,14.0]		13.3	[12.1,14.6]		10.8	[9.5,12.2]		10.8	[9.3,12.5]		0.8	-2.5**	0.0	-1.8
Female	15.1	[13.8,16.5]		15.5	[14.2,16.9]		12.1	[10.6,13.7]		12.4	[10.9,14.0]		0.4	-3.4**	0.3	-2.8**
Total	8,175			7,789			8,200			7,253						

¹Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions, p-value <0.05, **<0.01, ***<0.001. ²Children with don't know or missing information are included within the "other" category for place of delivery and assistance at delivery.

Appendix Table 9. Percentage of last-born children in the 2 years preceding the survey who started breastfeeding within 1 hour of birth, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000			2005			2010			2014			Difference ¹			
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014
Total	11.1	[9.9,12.4]		35.5	[33.3,37.9]		65.8	[63.3,68.2]		62.6	[60.0,65.0]		24.5***	30.3***	-3.2	51.5***
Place of residence																
Urban	13.1	[10.3,16.5]		36.9	[32.2,42.0]		64.4	[60.0,68.6]		50.6	[45.4,55.7]		23.9***	27.5***	-13.8***	37.5***
Rural	10.8	[9.6,12.1]		35.3	[32.8,37.9]		66.1	[63.3,68.8]		64.5	[61.7,67.2]		24.5***	30.8***	-1.6	53.7***
Region																
Phnom Penh	10.6	[6.0,18.1]		35.2	[30.0,40.7]		67.3	[59.4,74.3]		37.2	[30.2,44.8]		24.6***	32.1***	-30.1***	26.6***
Plain	11.0	[8.9,13.5]		36.1	[31.5,41.0]		62.9	[58.2,67.3]		68.2	[63.3,72.7]		25.1***	26.8***	5.3	57.2***
Great Lake	10.3	[8.5,12.4]		36.2	[32.7,39.9]		68.3	[64.1,72.2]		69.4	[65.5,73.0]		25.9***	32.1***	1.1	59.1***
Coastal	3.0	[1.2,7.3]		33.6	[27.4,40.4]		55.5	[48.2,62.5]		51.3	[42.2,60.3]		30.6***	21.9***	-4.2	48.3***
Plateau	18.5	[15.6,21.7]		33.7	[29.0,38.7]		72.0	[67.9,75.8]		55.0	[50.4,59.4]		15.2***	38.3***	-17.1***	36.5***
Wealth quintile																
Lowest	9.9	[7.9,12.4]	*	28.6	[24.1,33.5]	***	65.4	[60.5,70.0]		62.8	[57.0,68.3]		-18.7***	36.8***	-2.6	-52.9***
Second	9.2	[6.9,12.1]		31.1	[27.0,35.5]		64.4	[58.4,70.1]		62.2	[56.7,67.4]		-21.9***	33.3***	-2.2	-53.0***
Middle	9.8	[7.4,13.0]		41.1	[36.1,46.4]		68.0	[62.4,73.1]		71.0	[65.0,76.3]		-31.3***	26.8***	3.0	-61.1***
Fourth	13.5	[10.3,17.6]		41.3	[35.9,46.9]		69.0	[63.3,74.3]		64.1	[59.0,68.9]		27.8***	27.8***	-5.0	50.5***
Highest	15.3	[11.3,20.3]		40.5	[35.1,46.2]		62.0	[57.1,66.7]		52.8	[47.0,58.5]		25.2***	21.5***	-9.3*	37.5***
Mother's education																
No education	11.7	[9.6,14.1]		32.6	[28.1,37.5]	**	63.8	[58.0,69.2]		58.4	[50.9,65.6]		20.9***	31.2***	-5.3	46.7***
Primary	9.9	[8.3,11.8]		34.7	[32.0,37.5]		66.8	[63.5,70.0]		66.6	[63.0,70.0]		-24.8***	32.2***	-0.2	-56.7***
Secondary+	14.0	[10.5,18.4]		42.2	[36.5,48.3]		65.0	[60.5,69.2]		58.4	[53.5,62.1]		28.3***	22.7***	-6.6*	44.4***
Mother's age at birth																
<20	13.7	[9.9,18.8]		39.3	[32.1,47.0]		63.3	[55.9,70.0]		64.7	[58.0,71.0]		25.6***	23.9***	1.5	51.0***
20-34	11.1	[9.7,12.7]		35.1	[32.5,37.7]		66.5	[63.7,69.2]		62.3	[59.4,65.1]		24.0***	31.4***	-4.2*	51.2***
35-49	9.8	[7.3,13.2]		35.4	[31.1,39.9]		63.1	[56.6,69.2]		62.2	[53.6,70.1]		-25.5***	27.8***	-1.0	-52.4***
Birth order																
1	13.5	[10.7,17.0]		34.1	[29.7,38.8]		63.1	[59.1,66.8]	**	58.8	[55.2,62.3]		20.6***	28.9***	-4.3	45.2***
2-3	11.4	[9.4,13.7]		37.0	[33.6,40.6]		70.3	[66.9,73.5]		67.0	[63.5,70.3]		25.6***	33.3***	-3.4	55.6***
4-5	9.9	[7.6,12.8]		34.5	[29.1,40.3]		60.1	[53.7,66.2]		59.6	[52.8,66.1]		-24.7***	25.6***	-0.5	-49.7***
6+	9.7	[7.1,13.1]		35.3	[31.2,39.7]		61.5	[51.3,70.7]		60.6	[44.9,74.3]		-25.6***	26.2***	-0.9	-50.9***
Birth interval									*							
First Birth	13.6	[10.8,17.0]		34.1	[29.7,38.8]		62.6	[58.6,66.4]		58.7	[55.1,62.2]		20.5***	28.5***	-3.9	45.1***
<24	11.2	[8.1,15.3]		33.3	[27.6,39.6]		68.6	[61.0,75.3]		69.9	[61.9,76.8]		22.0***	35.3***	1.3	58.6***
24-47	10.6	[8.8,12.7]		38.0	[34.0,42.2]		66.9	[62.5,71.1]		65.5	[61.1,69.6]		27.4***	28.9***	-1.5	54.9***
48+	9.9	[7.6,12.8]		34.9	[30.9,39.1]		68.1	[63.6,72.3]		63.7	[59.2,68.0]		-25.0***	33.2***	-4.3	-53.8***
Place of delivery²									***							
Home, other	10.4	[9.2,11.8]	**	32.9	[30.3,35.6]	***	58.9	[54.6,63.0]		48.9	[40.4,57.4]		22.5***	26.0***	-10.0*	38.4***
Health facility	16.6	[12.0,22.7]		43.3	[38.4,48.3]		69.3	[66.5,72.0]		64.3	[61.8,66.7]		26.6***	26.1***	-5.0*	47.6***
Assistance at delivery²									***							
No one, other	9.9	[8.5,11.4]	*	29.8	[26.7,33.1]	***	57.4	[52.2,62.4]		45.0	[35.2,55.2]		-19.9***	27.6***	-12.4*	-35.2***
Doctor, nurse, midwife	13.4	[11.0,16.2]		41.6	[38.1,45.1]		68.0	[65.4,70.5]		63.9	[61.4,66.4]		28.2***	26.4***	-4.1*	50.5***
Child's sex									*							
Male	10.1	[8.6,11.9]		32.8	[29.9,35.8]	**	63.4	[60.2,66.5]		60.9	[57.4,64.2]		22.6***	30.7***	-2.5	50.8***
Female	12.1	[10.4,13.9]		38.4	[35.4,41.5]		68.3	[65.1,71.3]		64.2	[60.9,67.4]		26.4***	29.9***	-4.1	52.2***
T total	2,985			3,083			3,187			2,944						

¹ Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions, p-value * < 0.05, ** < 0.01, *** < 0.001. ² Children with don't know or missing information are included within the "other" category for place of delivery and assistance at delivery.

Appendix Table 10. Percentage of children age 12-23 months who are fully immunized, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000			2005			2010			2014			Difference ¹			
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014
Total	39.9	[35.9,44.0]		66.6	[63.2,69.8]		78.9	[75.7,81.7]		73.4	[70.2,76.5]		26.7***	12.2***	-5.4*	33.6***
Place of residence																
Urban	46.3	[37.1,55.7]		69.4	[62.4,75.5]	**	85.5	[80.7,89.2]	**	86.4	[82.1,89.9]	***	23.1***	16.1***	1.0	40.1***
Rural	39.0	[34.9,43.2]		66.2	[62.4,69.8]		77.5	[73.8,80.8]		71.2	[67.4,74.7]		27.2***	11.3***	-6.3*	32.2***
Region																
Phnom Penh	61.8	[45.0,76.2]	**	80.9	[66.6,90.0]	***	84.2	[75.0,90.4]	**	89.1	[80.0,94.4]	***	19.1*	3.3	5.0	27.3**
Plain	39.4	[32.2,47.0]		71.5	[65.1,77.1]		80.1	[73.5,85.4]		66.9	[60.0,73.2]		32.1***	8.6*	-13.2**	27.6***
Great Lake	41.9	[35.9,48.3]		63.6	[57.8,69.0]		81.9	[77.2,85.8]		81.4	[76.8,85.3]		21.6***	18.3***	-0.4	39.5***
Coastal	20.4	[9.4,38.7]		49.6	[40.3,58.9]		63.6	[51.9,74.0]		75.6	[65.3,83.6]		29.2**	14.1	12.0	55.2***
Plateau	37.6	[29.6,46.4]		61.8	[54.7,68.4]		72.7	[66.6,78.1]		62.1	[55.3,68.4]		24.2***	10.9*	-10.7*	24.4***
Wealth quintile																
Lowest	28.6	[22.5,35.6]	***	56.1	[48.9,63.0]	***	65.3	[57.9,72.1]	***	60.9	[53.6,67.7]	***	27.5***	9.2*	-4.4	32.3***
Second	34.7	[27.0,43.2]		65.8	[58.5,72.5]		77.7	[71.3,83.0]		65.4	[56.7,73.1]		31.2***	11.9**	-12.4**	30.7***
Middle	38.4	[29.1,48.7]		66.6	[60.9,71.9]		83.6	[77.7,88.2]		70.0	[60.1,78.2]		28.2***	17.0***	-13.7**	31.5***
Fourth	45.4	[36.5,54.7]		74.4	[62.5,83.5]		84.3	[76.5,89.8]		85.2	[78.6,90.0]		28.9***	9.9	0.9	39.7***
Highest	67.7	[56.6,77.1]		76.4	[65.6,84.6]		88.2	[82.5,92.2]		90.5	[86.2,93.6]		8.7	11.8**	2.3	22.8***
Mother's education																
No education	29.1	[23.2,35.9]	***	51.8	[42.9,60.5]	***	58.9	[50.2,67.1]	***	58.4	[49.6,66.6]	***	22.6***	7.1	-0.5	29.3***
Primary	41.4	[36.3,46.7]		67.6	[63.9,71.0]		80.1	[76.1,83.5]		69.7	[64.3,74.5]		26.2***	12.5***	-10.4***	28.3***
Secondary+	58.8	[47.2,69.6]		83.3	[76.4,88.5]		87.6	[83.1,91.1]		84.2	[80.2,87.5]		24.5***	4.4	-3.5	25.4***
Mother's age at birth																
<20	39.2	[29.4,50.1]		64.3	[48.5,77.4]		85.0	[79.2,89.4]	*	74.7	[63.6,83.3]		25.0***	20.7***	-10.3	35.4***
20-34	40.1	[35.7,44.7]		67.6	[63.3,71.5]		79.2	[75.6,82.4]		73.9	[70.1,77.4]		27.4***	11.7***	-5.3*	33.8***
35-49	39.4	[30.9,48.6]		64.1	[54.0,73.1]		71.7	[62.2,79.5]		68.1	[56.6,77.8]		24.7***	7.6	-3.6	28.7***
Birth Order																
1	40.6	[33.2,48.4]		69.7	[63.1,75.6]		83.8	[79.8,87.2]	***	75.9	[71.0,80.2]	*	29.1***	14.1***	-7.9**	35.3***
2-3	43.3	[36.9,49.9]		68.5	[62.3,74.1]		79.3	[75.1,83.0]		73.9	[69.0,78.3]		25.2***	10.9***	-5.4	30.6***
4-5	34.6	[28.1,41.6]		61.1	[52.6,69.0]		70.9	[60.4,79.6]		69.1	[59.0,77.6]		26.6***	9.8	-1.8	34.5***
6+	39.0	[31.6,46.8]		62.0	[53.2,70.1]		61.8	[46.8,74.8]		52.4	[36.0,68.3]		23.0***	-0.2	-9.4	13.4
Place of delivery²																
Home, other	37.3	[33.2,41.5]	***	65.2	[61.2,69.0]	***	67.8	[61.6,73.4]	***	50.2	[40.9,59.5]	***	28.0***	2.6	-17.6***	12.9**
Health facility	67.3	[52.6,79.2]		71.2	[61.3,79.3]		84.9	[82.0,87.5]		77.0	[73.5,80.0]		3.9	13.8***	-8.0***	9.7
Child's sex																
Male	42.6	[37.4,47.9]		69.0	[64.5,73.1]		77.2	[72.8,81.1]		73.9	[69.3,78.0]		26.4***	8.2**	-3.3	31.3***
Female	37.2	[32.1,42.5]		64.1	[59.8,68.2]		80.5	[76.6,84.0]		73.0	[68.6,76.9]		27.0***	16.4***	-7.6**	35.8***
Total	1,253			1,517			1,614			1,460						

¹Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions. p-values * <0.05 , ** <0.01 , *** <0.001 . ²Children with don't know or missing information are included within the "other" category for place of delivery.

Appendix Table 11. Among children under age 5 who experienced diarrhea in the 2 weeks before the survey, percentage for whom treatment was sought from a health facility or provider, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000			2005			2010			2014			Difference ¹			
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014
Total	21.6	[18.8,24.7]		41.1	[37.5,44.8]		58.9	[55.2,62.6]		55.5	[50.8,60.0]		19.5***	17.8***	-3.5	33.8***
Place of residence																
Urban	31.8	[22.7,42.5]	**	39.1	[29.1,50.0]		58.1	[49.3,66.5]		47.0	[39.1,55.0]	*	7.3	19.1**	-11.2	15.2*
Rural	20.3	[17.6,23.4]		41.4	[37.6,45.3]	**	59.0	[54.9,63.0]		56.9	[51.7,61.9]		21.0***	17.6***	-2.1	36.5***
Region																
Phnom Penh	26.4	[14.3,43.3]		44.1	[27.0,62.7]		59.5	[47.0,70.8]		57.5	[45.3,68.8]		17.7	15.4	-2.0	31.2**
Plain	23.7	[17.8,30.7]		48.5	[41.7,55.3]		64.5	[57.7,70.7]		55.2	[45.5,64.5]		24.8***	16.0***	-9.3	31.5***
Great Lake	17.0	[13.7,21.0]		36.1	[31.0,41.6]		53.8	[47.6,59.9]		52.5	[44.0,60.8]		19.1***	17.7***	-1.3	35.4***
Coastal	26.3	[17.8,37.1]		32.9	[18.9,50.9]		52.6	[41.2,63.6]		64.4	[51.1,75.9]		6.6	19.6*	11.9	38.1***
Plateau	22.9	[17.7,29.2]		31.2	[25.5,37.4]		57.8	[50.8,64.5]		57.4	[49.7,64.8]		8.2*	26.6***	-0.3	34.5***
Wealth quintile																
Lowest	20.5	[16.0,26.0]	*	35.0	[28.4,42.2]		53.3	[46.5,60.1]		62.0	[53.1,70.2]		14.5***	18.3***	8.7	41.5***
Second	15.7	[11.7,20.8]		40.5	[28.0,54.3]		59.4	[49.5,68.6]		53.6	[41.4,65.4]		24.7***	18.9**	-5.8	37.8***
Middle	20.6	[14.3,28.9]		47.5	[41.2,54.0]		67.3	[57.5,75.8]		57.8	[45.8,68.9]		26.9***	19.8***	-9.5	37.2***
Fourth	24.3	[17.9,32.1]		40.1	[29.0,52.4]		59.3	[47.6,70.0]		55.7	[42.3,68.3]		15.8**	19.1**	-3.6	31.4***
Highest	33.8	[21.6,48.6]		49.5	[37.7,61.4]		59.7	[47.4,70.8]		43.8	[33.6,54.6]		15.7*	10.2	-15.8*	10.1
Mother's education																
No education	20.1	[15.9,25.0]	*	37.3	[29.5,45.8]		55.5	[46.9,63.8]		58.8	[42.7,73.6]		17.2***	18.2***	3.3	38.8***
Primary	20.4	[16.9,24.5]		40.3	[35.0,45.8]		58.8	[54.2,63.3]		55.2	[48.4,61.9]		19.8***	18.6***	-3.6	34.8***
Secondary+	31.1	[22.3,41.5]		52.0	[38.9,64.8]		63.0	[55.2,70.1]		54.3	[47.0,61.5]		20.8**	11.0*	-8.6	23.2***
Mother's age at birth																
<20	26.6	[16.6,39.8]		39.2	[21.9,59.7]		61.2	[50.0,71.4]		62.9	[48.9,75.0]		12.6	22.0**	1.7	36.3***
20-34	21.1	[17.8,24.7]		42.7	[38.8,46.8]		60.1	[55.6,64.5]		55.0	[50.1,59.8]		21.7***	17.4***	-5.1	33.9***
35-49	21.1	[15.1,28.7]		35.2	[27.1,44.2]		48.8	[37.3,60.4]		45.0	[17.8,75.6]		14.1**	13.6*	-3.8	23.9*
Birth order																
1	24.6	[18.0,32.7]		43.4	[36.5,50.5]	**	61.5	[55.5,67.2]	*	56.6	[49.7,63.4]		18.8***	18.1***	-4.9	32.1***
2-3	23.3	[18.6,28.8]		46.3	[40.2,52.5]		61.6	[56.0,66.9]		57.7	[51.0,64.2]		23.0***	15.3***	-9.9	34.4***
4-5	17.2	[12.6,23.1]		35.1	[28.2,42.7]		53.3	[40.9,65.3]		46.6	[32.5,61.3]		17.8***	18.2**	-6.7	29.4***
6+	21.3	[15.5,28.5]		29.3	[20.5,40.1]		41.9	[28.6,56.5]		39.2	[12.6,74.3]		8.0	12.6	-2.8	17.9
Place of delivery²																
Home, other	21.2	[18.3,24.4]		38.6	[34.4,43.0]	**	55.5	[48.9,61.8]		56.6	[43.7,68.7]		17.4***	16.9***	1.2	35.5***
Health facility	25.7	[13.9,42.6]		51.3	[40.1,62.3]		62.0	[57.2,66.6]		55.2	[50.0,60.2]		25.5***	10.8*	-6.8	29.5***
Assistance at delivery²																
No one, other	18.5	[15.6,21.7]	**	37.0	[31.6,42.8]	**	50.1	[42.2,58.1]	**	49.1	[31.8,66.6]		18.6***	13.1**	-1.1	30.6***
Doctor, nurse, midwife	28.0	[22.4,34.3]		47.5	[40.9,54.2]	**	62.8	[58.6,66.7]		56.4	[51.3,61.3]		19.5***	15.3***	-6.4	28.4***
Child's current age																
<6 mo	13.5	[7.7,22.4]	**	29.1	[22.5,36.8]	**	51.5	[36.5,66.2]	*	43.6	[30.5,57.7]		15.7*	22.4**	-7.8	30.2***
6-11 mo	24.7	[18.6,32.0]		40.8	[31.7,50.7]		64.4	[55.3,72.6]		58.1	[45.6,69.5]		16.2**	23.6***	-6.3	33.4***
1 year	28.7	[23.1,35.0]		50.3	[43.2,57.5]		66.6	[60.1,72.5]		57.4	[48.6,65.8]		21.6**	16.2***	-9.1	28.8***
2 years	23.8	[18.1,30.7]		41.3	[32.2,51.0]		58.3	[43.5,65.6]		54.8	[43.5,65.6]		17.5***	17.1**	-3.6	31.0***
3 years	13.4	[8.7,20.1]		28.7	[19.7,39.7]		49.5	[38.3,60.7]		57.3	[40.8,72.4]		15.3***	20.7**	7.9	43.9***
4 years	16.8	[11.1,24.5]		41.8	[34.4,49.6]		47.4	[35.3,59.7]		56.7	[40.4,71.6]		25.0***	5.6	9.3	39.9***
Child's sex																
Male	21.4	[17.9,25.3]		42.6	[37.4,47.9]		57.1	[51.6,62.4]		53.1	[46.8,59.3]		21.2***	14.5***	-4.0	31.7***
Female	21.9	[17.7,26.9]		39.3	[34.3,44.5]		61.2	[55.7,66.4]		58.1	[51.2,64.6]		17.4***	21.9***	-3.1	36.1***
Total	1,385			1,420			1,161			902						

¹ Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions. p-values * < 0.05, ** < 0.01, *** < 0.001. ² Children with don't know or missing information are included within the "other" category for place of delivery and assistance at delivery.

Appendix Table 12. Among children under age 5 who experienced diarrhea in the 2 weeks before the survey, percentage given ORT or increased fluids, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014 DHS

	2000			2005			2010			2014			Difference ¹				
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014	
Total	61.5	[58.2,64.6]		49.1	[44.9,53.3]		52.6	[48.6,56.6]		57.0	[51.7,62.2]		-12.4***	3.5	4.4	4.4	
Place of residence																	
Urban	64.6	[55.6,72.8]		44.5	[38.1,51.1]		51.8	[42.8,60.8]		59.1	[48.4,69.1]		-20.1***	7.3	7.3	5.5	
Rural	61.1	[57.8,64.3]		49.7	[45.1,54.3]		52.7	[48.4,57.1]		56.7	[50.8,62.4]		-11.3***	3.0	4.0	4.4	
Region																	
Phnom Penh	71.7	[59.9,81.0]		27.0	[14.0,45.6]	**	52.1	[37.0,66.9]		69.2	[57.0,79.3]		-44.7***	25.1*	-17.1	2.4	
Plain	56.6	[49.6,63.3]		48.0	[39.2,57.0]		50.3	[43.4,57.1]		52.3	[41.1,63.3]		-8.6	2.3	2.0	4.3	
Great Lake	68.6	[64.1,72.9]		52.6	[47.1,58.1]		57.4	[50.4,64.1]		64.7	[55.6,72.9]		-16.0***	4.8	7.3	3.9	
Coastal	60.4	[50.8,69.4]		45.1	[33.6,57.1]		42.4	[33.1,52.3]		51.9	[36.7,66.7]		-15.4*	-2.7	9.5	-8.6	
Plateau	47.5	[39.8,55.4]		57.6	[52.1,62.9]		52.4	[44.7,60.0]		45.8	[36.8,55.2]		10.1*	-5.1	6.6	-1.7	
Wealth																	
Lowest	59.3	[52.8,65.5]		54.1	[46.9,61.2]	***	54.2	[45.2,63.0]		56.3	[45.4,66.7]		-5.2	0.1	2.1	3.0	
Second	62.0	[55.9,67.8]		55.5	[44.7,65.8]		45.5	[35.9,55.5]		53.5	[41.7,64.9]		-6.5	-10.0	8.0	-8.6	
Middle	58.5	[49.2,67.3]		48.5	[39.1,58.1]		60.1	[50.2,69.2]		57.1	[44.4,68.9]		-10.0	11.5*	3.0	1.4	
Fourth	63.3	[54.1,71.6]		43.2	[35.1,51.6]		52.8	[41.6,63.7]		57.6	[39.4,73.9]		-20.1***	9.6	4.8	5.7	
Highest	67.2	[55.9,76.8]		32.1	[22.5,43.4]		50.2	[38.0,62.3]		61.4	[49.6,72.1]		-35.1***	18.1*	11.3	5.7	
Mother's education																	
No education	59.4	[54.1,64.6]		48.6	[39.8,57.4]		52.0	[44.1,59.7]		61.3	[44.3,75.9]		-10.9*	3.4	9.3	1.8	
Primary	62.6	[57.9,67.1]		51.4	[45.6,57.3]		53.8	[48.5,58.9]		54.9	[48.0,61.7]		-11.2**	2.3	1.2	7.6	
Secondary+	61.7	[50.7,71.5]		40.1	[28.8,52.4]		49.7	[41.7,57.7]		58.6	[50.6,66.2]		-21.6**	9.7	8.9	3.1	
Mother's age at birth																	
<20	60.1	[45.6,73.0]		36.7	[23.5,52.3]		60.2	[45.7,73.2]		62.2	[50.1,72.9]		-23.4**	23.5**	1.9	2.1	
20-34	62.7	[58.7,66.5]		50.7	[45.6,55.7]		50.9	[46.4,55.3]		57.5	[52.0,62.9]		-12.0***	0.2	6.6	5.2	
35-49	57.7	[49.7,65.3]		50.1	[41.6,58.6]		56.2	[45.9,65.9]		38.9	[18.3,64.5]		-7.6	6.1	17.3	18.8	
Birth order																	
1	59.5	[51.5,67.1]		44.4	[35.5,53.7]		53.4	[46.1,60.7]		61.3	[54.2,67.9]		-15.1**	9.0	7.8	1.8	
2-3	62.9	[56.9,68.5]		49.4	[42.8,56.1]		54.2	[48.6,59.6]		54.5	[47.1,61.7]		-13.4***	4.7	0.3	8.4	
4-5	60.3	[54.4,65.9]		54.9	[45.7,63.7]		48.2	[36.4,60.1]		57.8	[42.8,71.6]		-5.4	-6.7	9.7	-2.4	
6+	62.1	[54.1,69.4]		49.9	[40.7,59.1]		48.3	[34.5,62.3]		32.6	[8.0,73.1]		-12.2*	-1.6	15.7	-29.4*	
Place of delivery²																	
Home, other	61.0	[57.6,64.4]		51.4	[46.3,56.4]	*	51.8	[46.1,57.4]		42.7	[26.7,60.4]		-9.6***	0.4	9.1	18.3**	
Health facility	65.5	[52.9,76.3]		39.9	[29.4,51.4]		53.4	[47.8,58.9]		60.1	[54.5,65.5]		-25.6***	13.5**	6.7	5.4	
Child's current age																	
<6 mo	32.8	[24.4,42.4]		23.1	[17.1,30.3]	***	42.5	[27.6,58.9]		39.1	[22.2,59.1]		-9.7	19.4*	3.4	6.3	
6-11 mo	52.4	[44.8,59.9]		44.8	[33.2,57.1]		50.0	[40.9,59.1]		61.7	[50.6,71.7]		-7.6	5.2	11.7	9.3	
1 years	65.7	[59.6,71.4]		60.4	[51.5,68.6]		55.5	[47.5,63.1]		57.7	[49.1,65.9]		-5.4	-4.9	2.2	-8.1	
2 years	70.4	[63.5,76.4]		53.6	[45.1,61.9]		58.3	[48.1,67.8]		60.1	[49.2,70.1]		-16.8***	4.7	1.8	10.3	
3 years	64.7	[57.0,71.6]		44.6	[30.1,60.0]		48.7	[37.9,59.6]		57.7	[38.5,74.8]		-20.1**	4.1	9.0	7.0	
4 years	68.5	[59.8,76.0]		44.0	[36.9,51.3]		52.4	[39.1,65.4]		59.0	[40.5,75.2]		-24.5***	8.5	6.6	9.5	
Child's sex																	
Male	59.9	[55.6,64.1]		50.1	[44.5,55.8]		51.4	[46.0,56.9]		56.2	[49.7,62.4]		-9.8**	1.3	4.7	3.8	
Female	63.3	[58.1,68.1]		47.8	[41.4,54.4]		54.1	[48.3,59.8]		58.0	[50.9,64.8]		-15.4***	6.3	3.9	5.2	
Total	1,385			1,420			1,161			902							

¹ Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions. p-values * <0.05 , ** <0.01 , *** <0.001 . ² Children with don't know or missing information are included within the "other" category for place of delivery.

Appendix Table 13. Among children under age 5, percentage who are stunted, according to background characteristics, Cambodia 2000, 2005, 2010, and 2014
DHS

	2000			2005			2010			2014			Difference ¹			
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2000-2005	2005-2010	2010-2014	2000-2014
Total	49.6	[47.3,51.9]		42.4	[40.0,44.7]		39.1	[36.9,41.4]		31.9	[30.0,33.9]		-7.2***	-3.3	-7.2***	-17.7***
Place of residence																
Urban	41.9	[35.9,48.2]	**	34.3	[27.3,42.2]	*	26.9	[23.5,30.6]		23.1	[19.8,26.8]		-7.6	-7.4	-3.8	-18.8***
Rural	50.9	[48.6,53.2]		43.6	[41.1,46.2]		41.3	[38.8,43.9]		33.4	[31.3,35.6]		-7.2***	-2.3	-7.9***	-17.5***
Region																
Phnom Penh	30.8	[24.6,37.7]	***	24.7	[11.8,44.6]	**	25.1	[19.4,31.7]		16.4	[12.0,22.0]		-6.0	0.3	-8.6*	-14.4***
Plain	53.1	[48.5,57.6]		39.5	[35.1,44.1]		37.6	[33.2,42.1]		31.3	[28.2,34.6]		-13.6***	-1.9	-6.2*	-21.7***
Great Lake	48.0	[44.8,51.3]		49.6	[46.1,53.1]		40.1	[36.7,43.5]		33.4	[29.7,37.2]		1.6	-9.6***	-6.7*	-14.7***
Coastal	48.0	[38.7,57.5]		38.2	[33.0,43.6]		41.6	[35.2,48.3]		25.6	[21.2,30.5]		-9.9*	3.4	-16.0***	-22.4***
Plateau	51.9	[47.9,56.0]		45.2	[41.6,49.0]		47.7	[43.3,52.1]		39.9	[35.5,44.5]		-6.7*	2.4	-7.8*	-12.1***
Wealth																
Lowest	57.9	[53.4,62.3]	***	52.3	[48.0,56.7]	***	49.4	[44.8,54.0]		41.2	[36.9,45.6]		-5.6*	-3.0	-8.2**	-16.7***
Second	53.2	[48.9,57.5]		47.8	[43.7,52.0]		44.1	[39.4,48.9]		37.3	[33.2,41.6]		-5.4	-3.7	-6.8*	-15.9***
Middle	47.5	[41.9,53.0]		44.3	[38.5,50.3]		38.7	[33.6,44.0]		32.2	[27.6,37.3]		-3.1	-5.7	-6.4	-15.2***
Fourth	48.8	[42.9,54.7]		37.8	[31.4,44.6]		34.0	[29.4,38.8]		26.9	[23.2,31.0]		-11.0*	-3.8	-7.0*	-21.9***
Highest	31.6	[26.4,37.3]		22.9	[17.6,29.4]		21.9	[18.2,26.2]		18.4	[15.3,21.9]		-8.6*	-1.0	-3.5	-13.2***
Mother's education																
No education	55.6	[51.3,59.8]	***	52.0	[47.3,56.8]	***	46.3	[41.6,51.2]		39.0	[33.4,44.9]		-3.5	-5.7	-7.3*	-16.5***
Primary	48.3	[45.4,51.3]		43.5	[40.4,46.5]		40.2	[37.4,43.1]		33.9	[31.3,36.6]		-4.8*	-3.2	-6.3**	-14.4***
Secondary+	41.0	[34.7,47.7]		25.4	[20.5,31.1]		30.9	[27.0,34.9]		25.6	[22.7,28.7]		-15.6***	5.4	-5.2*	-15.4***
Mother's age at birth																
<20	56.4	[48.6,63.9]		40.3	[32.7,48.4]		43.3	[37.0,49.7]		33.4	[27.6,39.6]		-16.1***	2.9	-9.9*	-23.1***
20-34*	48.7	[46.1,51.3]		41.9	[39.1,44.8]		38.6	[36.2,41.2]		31.3	[29.2,33.4]		-6.8***	-3.2	-7.4***	-17.4***
35-49	49.7	[44.5,55.0]		45.4	[39.8,51.0]		38.6	[33.2,44.4]		36.0	[30.3,42.1]		-4.4	-6.7	-2.7	-13.8***
Mother smokes cigarettes																
No	48.5	[46.2,50.8]	**	41.2	[38.8,43.7]	***	38.4	[36.1,40.7]		31.3	[29.4,33.3]		-7.3***	-2.8	-7.1***	-17.1***
Yes	64.5	[53.3,74.3]		62.6	[55.2,69.4]		60.2	[49.0,70.5]		51.9	[40.2,63.5]		-2.0	-2.3	-8.3	-12.6*
Birth interval																
First Birth	46.6	[42.2,51.1]	***	35.3	[30.5,40.4]	***	36.0	[32.8,39.4]		30.1	[27.2,33.2]		-11.3***	0.8	-5.9*	-16.5***
<24	59.1	[53.9,64.2]		52.4	[45.8,58.9]		45.4	[39.2,51.7]		37.4	[30.5,44.8]		-6.7	-7.0	-8.0	-21.8***
24-47	51.8	[48.7,54.9]		45.9	[41.8,50.1]		43.3	[39.5,47.1]		34.0	[30.7,37.5]		-5.9*	-2.7	-9.2***	-17.8***
48+	40.1	[35.3,45.2]		39.7	[35.7,43.8]		35.9	[31.8,40.2]		30.9	[27.6,34.4]		-0.4	-3.8	-5.0	-9.2**
Place of delivery²																
Home, other*	51.4	[48.9,53.8]	***	45.8	[43.5,48.2]	***	48.4	[45.0,51.9]		45.4	[40.3,50.6]		-5.5***	2.6	-3.0	-6.0*
Health facility	35.1	[29.3,41.4]		29.6	[22.2,38.3]		30.7	[28.1,33.5]		29.4	[27.4,31.6]		-5.5	1.1	-1.3	-5.7*
Assistance at delivery²																
No one, other	53.0	[50.1,55.9]	***	50.0	[47.3,52.7]	***	52.8	[48.9,56.6]		45.7	[39.9,51.6]		-3.0	2.8	-7.1*	-7.3*
Doctor, nurse, midwife	42.7	[39.1,46.5]		33.0	[29.0,37.1]		33.5	[31.0,36.0]		30.4	[28.4,32.4]		-9.8***	0.5	-3.1	-12.4***

Continued...

Appendix Table 13—Continued

	2000			2005			2010			2014			Difference ¹		
	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	%	95% CI	p-value	2005-2010	2010-2014	2000-2014
Child's current age															
<6 mo	24.4	[19.3,30.4]	***	14.7	[10.1,21.1]	***	10.5	[6.7,16.2]	***	16.3	[12.2,21.5]	***	-4.2	5.8	-8.1*
6-11 mo	31.1	[25.3,37.5]		17.5	[13.4,22.5]		19.6	[15.4,24.5]		15.7	[11.4,21.2]		2.1	-3.9	-15.4***
1 years	45.3	[40.2,50.5]		42.7	[37.1,48.4]		39.5	[35.5,43.6]		31.0	[27.3,35.0]		-3.2	-8.4**	-14.3***
2 years	58.6	[54.1,63.0]		47.8	[42.7,52.9]		48.0	[43.4,52.6]		37.8	[33.8,42.1]		0.2	-10.1**	-20.8***
3 years	59.0	[54.7,63.3]		53.4	[48.3,58.5]		43.0	[38.5,47.5]		39.0	[34.3,44.0]		-5.6	-10.5***	-20.0***
4 years	57.1	[52.5,61.6]		50.6	[45.7,55.5]		48.2	[43.8,52.7]		36.5	[32.5,40.7]		-2.4	-11.8***	-20.6***
Child's sex															
Male	49.7	[46.5,52.8]	**	45.2	[42.0,48.4]	**	40.9	[38.1,43.9]	*	32.4	[29.8,35.1]		-4.3	-8.5***	-17.3***
Female	49.6	[46.7,52.4]		39.7	[36.9,42.5]		37.1	[34.3,40.0]		31.4	[28.9,34.1]		-2.6	-5.7**	-18.2***
Total	3,263			3,347			3,718			4,397					

¹Percentage point difference between two surveys and between 2000 and 2014 with significant tests for the difference in proportions. p-values * <0.05 , ** <0.01 , *** <0.001 . ²Children with don't know or missing information are included within the "other" category for place of delivery and assistance at delivery.

REFERENCES

- Adjiwanou, V., and T. LeGrand. 2013. "Does Antenatal Care Matter in the Use of Skilled Birth Attendance in Rural Africa: A Multi-Country Analysis." *Social Science & Medicine* 86:26-34.
- Alexander, G. R., and C. C. Korenbrot. 1995. "The Role of Prenatal Care in Preventing Low Birth Weight." *The Future of Children*: 103-120.
- Assaf, S., M. T. Kothari, and T. Pullum. 2015. *An Assessment of the Quality of DHS Anthropometric Data, 2005-2014*. DHS Methodological Reports No. 16. Rockville, Maryland, USA: ICF International. <http://dhsprogram.com/pubs/pdf/MR16/MR16.pdf>.
- Berkat, S., and R. Sutan. 2014. "The Effect of Early Initiation of Breastfeeding on Neonatal Mortality among Low Birth Weight in Aceh Province, Indonesia: An Unmatched Case Control Study." *Advances in Epidemiology* 2014.
- Betran, A., M. Torloni, J. Zhang, and A. Gülmezoglu. 2016. "Who Statement on Caesarean Section Rates." *BJOG: An International Journal of Obstetrics & Gynaecology* 123(5):667-670.
- Cambodia (MOH), T.M.o.H.K.o., and WHO. 2014. *Cambodia's Achievements in Bringing Down Death Rates of Mothers and Children Acknowledged Internationally*
- Chan Soeung, S., J. Grundy, R. Duncan, R. Thor, and J.B. Bilous. 2012. "From Reaching Every District to Reaching Every Community: Analysis and Response to the Challenge of Equity in Immunization in Cambodia." *Health Policy and Planning* 28(5):526-535.
- Dingle, A., C. Goodman, and T. Powell-Jackson. 2013. "A Decade of Improvements in Equity of Access to Reproductive and Maternal Health Services in Cambodia, 2000–2010." *International Journal for Equity in Health* 12(1):51.
- Flores, G., P. Ir, C. R. Men, O. O'Donnell, and E. Van Doorslaer. 2013. "Financial Protection of Patients through Compensation of Providers: The Impact of Health Equity Funds in Cambodia." *Journal of Health Economics* 32(6):1180-1193.
- Forsberg, B. C., C. T. Sreeramareddy, and Y.-P. Low. 2017. "Slow Progress in Diarrhea Case Management in Low and Middle Income Countries: Evidence from Cross-Sectional National Surveys, 1985–2012." *BMC Pediatrics* 17(1):83.
- Ikeda, N., Y. Irie, and K. Shibuya. 2013. "Determinants of Reduced Child Stunting in Cambodia: Analysis of Pooled Data from Three Demographic and Health Surveys." *Bulletin of the World Health Organization* 91(5):341-349.
- Ir, P., D. Horemans, N. Souk, and W. Van Damme. 2010. "Using Targeted Vouchers and Health Equity Funds to Improve Access to Skilled Birth Attendants for Poor Women: A Case Study in Three Rural Health Districts in Cambodia." *BMC Pregnancy and Childbirth* 10(1):1.

- Kerber, K. J., J. E. de Graft-Johnson, Z. A. Bhutta, P. Okong, A. Starrs, and J. E. Lawn. 2007. "Continuum of Care for Maternal, Newborn, and Child Health: From Slogan to Service Delivery." *The Lancet* 370(9595):1358-1369.
- National Institute of Public Health/Cambodia, National Institute of Statistics/Cambodia, and ORC Macro. 2006. *Cambodia Demographic and Health Survey 2005*. Phnom Penh, Cambodia: National Institute of Public Health/Cambodia, National Institute of Statistics/Cambodia, and ORC Macro. <http://dhsprogram.com/pubs/pdf/FR185/FR185.pdf>.
- National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ICF International. 2015. *Cambodia Demographic and Health Survey 2014*. Phnom Penh, Cambodia: National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ICF International. <http://dhsprogram.com/pubs/pdf/FR312/FR312.pdf>.
- National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ICF Macro. 2011. *Cambodia Demographic and Health Survey 2010*. Phnom Penh, Cambodia: National Institute of Statistics, Directorate General for Health, and ICF Macro. <http://dhsprogram.com/pubs/pdf/FR249/FR249.pdf>.
- National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ORC Macro. 2001. *Cambodia Demographic and Health Survey 2000*. Phnom Penh, Cambodia: National Institute of Statistics/Cambodia, Directorate General for Health/Cambodia, and ORC Macro. <http://dhsprogram.com/pubs/pdf/FR124/FR124.pdf>.
- Prak, S., M. I. Dahl, S. Oeurn, J. Conkle, A. Wise, and A. Laillou. 2014. "Breastfeeding Trends in Cambodia, and the Increased Use of Breast-Milk Substitute—Why Is It a Danger?" *Nutrients* 6(7):2920-2930.
- Rikimaru, T. 2015. "What Are the Current Situations and the Challenges of Maternal and Child Malnutrition in Asia?" *Journal of nutritional science and vitaminology* 61(Supplement):S63-S65.
- UNGA. 2015. *Transforming Our World: The 2030 Agenda for Sustainable Development*.
- Van de Poel, E., G. Flores, P. Ir, and E. Van Doorslaer. 2014. "Can Vouchers Deliver? An Evaluation of Subsidies for Maternal Health Care in Cambodia." *Bulletin of the World Health Organization* 92(5):331-339.
- Wang, W., and R. Hong. 2015. "Levels and Determinants of Continuum of Care for Maternal and Newborn Health in Cambodia-Evidence from a Population-Based Survey." *BMC Pregnancy and Childbirth* 15(1):62.
- WHO. n.d. *Who in the Western Pacific Region*. WHO. http://www.wpro.who.int/about/administration_structure/dhs/story_cambodia_reduces_maternal_mortality/en/.
- Zanello, G., C. Srinivasan, and B. Shankar. 2016. "What Explains Cambodia's Success in Reducing Child Stunting-2000-2014?" *PloS One* 11(9):e0162668.